

THE PILLOWCASE AND TRACELESS REPRESENTATIONS OF KNOT GROUPS II: A LAGRANGIAN-FLOER THEORY IN THE PILLOWCASE

MATTHEW HEDDEN, CHRISTOPHER M. HERALD, AND PAUL KIRK

ABSTRACT. We define an elementary relatively $\mathbb{Z}/4$ graded Lagrangian-Floer chain complex for restricted immersions of compact 1-manifolds into the pillowcase, and apply it to the intersection diagram obtained by taking traceless $SU(2)$ character varieties of 2-tangle decompositions of knots. Calculations for torus knots are explained in terms of pictures in the punctured plane. The relation to the reduced instanton homology of knots is explored.

1. INTRODUCTION

In [21], Kronheimer and Mrowka introduced a powerful invariant of a knot or link in a 3-manifold $L \subset X$ called *singular instanton knot homology*. Denoted $I^\natural(Y, L)$, their invariant is defined in the context of gauge theory. Roughly, the theory associates to L a chain group $C^\natural(X, L)$ generated by flat $SO(3)$ connections on $X \setminus L$ which have a prescribed singularity near L . This group is endowed with a differential that counts anti-self-dual instantons on $X \times \mathbb{R}$ which limit to given flat connections on the ends. Singular instanton knot homology has an important computational tool called the *skein exact triangle*. This is a long exact sequence relating the homology groups of links which agree outside of a small 3-ball, where they differ in a simple way. Iterated application of the exact triangle using a collection of 3-balls leads to a spectral sequence which converges to $I^\natural(X, L)$. This spectral sequence, when applied to a collection of 3-balls containing all the crossings of a diagram for a link in the 3-sphere, has E_2 page isomorphic to the well-known combinatorial knot invariant Khovanov homology [23]. The existence of this spectral sequence, together with a non-triviality result for $I^\natural(S^3, L)$ coming from its relation to another knot invariant (sutured instanton knot homology) [22], allowed Kronheimer and Mrowka to prove the striking result that Khovanov homology detects the unknot. Despite this triumph, $I^\natural(X, L)$ remains rather mysterious. This is due in large part to the fact that computations are extremely scarce. Initially, the only route for computation was through the aforementioned spectral sequence, but, aside from the instances where it collapses for simple reasons at Khovanov homology, little headway has been made in this direction (though see [24, 27] for more sophisticated computations using the spectral sequence).

Motivated by a desire for a more explicit understanding of the singular instanton chain complexes, we began a project in [18] which aims to make concrete direct calculations of $I^\natural(X, L)$. This is not so easy, and a serious initial sticking point arises from the fact that the flat connections which generate $C^\natural(X, L)$ are never isolated. Indeed, aside from the case

2010 *Mathematics Subject Classification.* Primary 57M27, 57R58, 57M25 ; Secondary 81T13.

Key words and phrases. pillowcase, holonomy perturbation, instanton, Floer homology, character variety, two bridge knot, torus knot.

of the unknot in S^3 , the spaces of flat connections studied by the theory are always positive dimensional varieties. For this reason, one needs to perturb the Chern-Simons functional that gives rise to $I^\natural(X, L)$ through its Morse homology. The robust holonomy perturbations used to set up the general theory destroy the concrete algebraic interpretation of the generating set for $C^\natural(X, K)$ in terms of certain traceless representations of the fundamental group of $X \setminus L$, and the goal of our first paper was to retain such an interpretation by way of explicit local, and in some sense minimal, holonomy perturbations. The main idea from [18] was to pick a particular distinguished 3-ball in X which intersects L in a trivial 2-stranded tangle. We then performed an explicit holonomy perturbation to the Chern-Simons functional in the neighborhood of a curve living in this ball. Using this perturbation allowed us, in a variety of examples, to perform computations of singular instanton chain groups for many knots (e.g., many torus knots) which implied that the spectral sequence from Khovanov homology necessarily had large rank higher differentials. In many of these cases we produced perfect complexes, i.e., complexes with trivial differential, so that our computation determined the singular instanton homology despite the fact that the spectral sequence from Khovanov homology was not understood.

The key perspective for the computations of [18] is that the generators of the singular instanton chain groups can be interpreted as the intersections of two immersed 1-manifolds in a 2-dimensional orbifold, *the pillowcase*, which arises as the quotient of the torus by the hyperelliptic involution (see Section 3 for more details). This perspective results from the observation that the choice of trivial 2-stranded tangle $(D, U) \subset (X, L)$ where we perform the perturbation results in a decomposition of the link

$$(X, L) = (X \setminus D, L \setminus U) \cup_{(S^2, \{4 \text{ points}\})} (D, U)$$

We let $(Y, T) := (X \setminus D, L \setminus U)$ denote the complementary tangle. Now $C^\natural(X, L)$ is generated by certain conjugacy classes of perturbed traceless representations $\rho : \pi_1(X \setminus L) \rightarrow SU(2)$. Using the above decomposition, they can be viewed as the intersection of the restrictions

$$\begin{array}{ccc} R(Y, T) & & R_\pi^\natural(D, U) \\ & \searrow \quad \swarrow & \\ & R(S^2, \{4 \text{ points}\}) \simeq P \end{array}$$

where $R(Y, T)$ denotes the traceless $SU(2)$ character variety of the complementary tangle, $R_\pi^\natural(D, U)$ denotes the perturbed traceless character variety of the trivial tangle (suitably summed with the Hopf link to allow for a non-trivial bundle), and $R(S^2, \{4 \text{ points}\})$ is the traceless character variety of the 4-punctured 2-sphere where the tangles intersect. The latter variety is isomorphic to the pillowcase, which we denote by P [18, Proposition 3.1]. In [18, Theorem 1] we calculated the restriction map $L_0 : R_\pi^\natural(D, U) \rightarrow P$, showing that for certain perturbation data its image was an immersed circle with exactly one double point, a “figure eight”. Provided that the image of the restriction $L_1 : R(Y, T) \rightarrow P$ is an immersed 1-manifold transverse to this figure eight, there is a bijection between generators for $C^\natural(X, L)$

and intersections of the images of L_0 and L_1 :

$$C^\natural(X, L) = \bigoplus_{x \in \{\text{Image } L_0 \cap \text{Image } L_1\}} \mathbb{Z}/2\langle x \rangle$$

This perspective allows for the computation of $C^\natural(X, L)$ for an arbitrary 2-bridge or torus knot and for certain pretzel knots (see [18, Sections 10 and 11] and [17]).

Though it made progress towards our goal of making the singular instanton complexes more computable, the approach of [18] had two serious drawbacks. The first is that it was not clear whether, given a link (X, L) , a trivial tangle $(D, U) \subset (X, L)$ can be found for which $L_1 : R(Y, T) \rightarrow P$ is an immersed 1-manifold transverse to the image of L_0 . The second is that even when such a tangle can be found, we had no way to compute the instanton differential on the resulting chain group. The purpose of the present article is to address this second issue.

A hint towards a possible understanding of the differential is gleaned by viewing our setup through the lens of an ever growing body of conjectured or established relationships between gauge theoretic and symplectically defined Floer theories (e.g., [2, 3, 32, 12, 34, 35, 8] discuss relationships between Yang-Mills gauge theory and symplectic invariants). These relationships are often described as “Atiyah-Floer Conjectures”, and our description of $C^\natural(X, L)$ suggests looking for a differential on the instanton complex in terms of the symplectic geometry of the pillowcase. In fact such a differential exists, and in the first half of this paper we introduce an elementary $\mathbb{Z}/4$ relatively graded Lagrangian-Floer type chain complex for appropriate immersions of compact 1-manifolds (*restricted immersed Lagrangians*) into the pillowcase. That is, we define a complex generated by intersections of the images of immersed 1-manifolds, whose boundary operator is defined by counting the analog of holomorphic disks connecting them (in this low-dimensional setting, it will suffice to count orientation preserving immersions of disks into the pillowcase connecting intersections of L_0 and L_1). In pursuit of our chain complexes, we draw liberally from the foundational work of Abouzaid [1] on Floer homology for immersed Lagrangians in Riemann surfaces and de Silva-Robbin-Salamon [29] for combinatorial and homotopy-theoretic aspects of Lagrangian Floer homology in this setting. Our work here can be viewed both as generalization and specialization of the existing literature, and our primary contribution is clarifying invariance proofs and properties of immersed Lagrangian Floer homology in the $\mathbb{Z}/4$ graded setting, and when some of the Lagrangians are immersed arcs (as opposed to circles). The main result towards this end is Theorem 4.1, which can be paraphrased as follows.

Theorem 1. *Let (L_0, L_1) be a pair of restricted immersed 1-manifolds in the pillowcase such that at least one of L_i consists only of circles. Then there is a well-defined Floer homology group $HF(L_0, L_1)$ whose relatively $\mathbb{Z}/4$ graded isomorphism type depends only on the free homotopy type of the pair (L_0, L_1) .*

The second half of the article applies this construction to the situation described above, when $L_0 : R_\pi^\natural(D, U) \rightarrow P$ is the restriction map from the perturbed traceless $SU(2)$ character variety of the trivial tangle and $L_1 : R_\pi(Y, T) \rightarrow P$ is the restriction of the perturbed traceless character variety of a tangle T in a homology 3-ball Y (e.g., the complementary tangle to an embedded trivial tangle in a pair (X, L) as above). In favorable circumstances, such as when (Y, T) is a certain tangle naturally associated to a 2-bridge or torus knot, the map

$L_1 : R_\pi(Y, T) \rightarrow P$ is a restricted Lagrangian without perturbations, so that the chain complex $C^\natural(Y, T, \pi) := CF(L_0, L_1)$ and its homology $H^\natural(Y, T, \pi)$, which we refer to as the **pillowcase homology** of (Y, T) , are defined. We then calculate the pillowcase homology for a number of examples, and show that it agrees with the singular instanton homology in cases where the latter is known (e.g., 2-bridge knots and many torus knots). More generally, our computations of pillowcase homology agree with conjectures for $I^\natural(S^3, K)$ in many other cases. We make the following Atiyah-Floer type conjecture:

Conjecture. *Given a knot K in a homology sphere X , there exists a 2-tangle decomposition $(X, K) = (Y, T) \cup (D, U)$ with (D, U) a trivial 2-tangle, and arbitrarily small perturbations π , so that $L_1 : R_\pi(Y, T) \rightarrow P$ is a restricted immersed 1-manifold for which the resulting pillowcase homology $H^\natural(Y, T, \pi)$ is isomorphic to $I^\natural(X, K)$ as a $\mathbb{Z}/4$ relatively graded group.*

We should note that the holonomy perturbations π that we use to make the traceless character varieties regular and transverse are compatible with the perturbations used to make the moduli spaces regular in the construction of $I^\natural(S^3, K)$. We should also note that while we have in some sense dealt with the second drawback from our first paper, in the sense that we have constructed a differential, the present results still leave us quite far from achieving our goal of computing $I^\natural(X, K)$. Indeed, we do not yet know how to construct the general perturbations necessary to even define the pillowcase homology (nor do we know how to pick the embedded trivial tangle (D, U)). Moreover, even in the cases that we can find perturbations which make the pillowcase homology well-defined, it is not true that any such perturbations will yield a complex whose homology agrees with $I^\natural(X, K)$.

We hope to remedy these concerns for the case of links in the 3-sphere in a subsequent article, which will develop a spectral sequence from Khovanov homology to pillowcase homology. This will rely on picking a particular trivial tangle with which to apply our construction, and then iterating a skein exact triangle satisfied by pillowcase homology in a similar manner to Kronheimer and Mrowka's construction. This will rely on picking diagrammatically defined perturbation curves for the diagram of the complementary tangle (Y, T) . Provided that we can construct this spectral sequence, it will have the advantage of providing an algebraic route to proving not only that pillowcase homology can be defined for arbitrary links in the 3-sphere, but also for showing that it is an invariant of the isotopy type of the link (in reality, we will prove the stronger result that the quasi-isomorphism type of a certain twisted complex built from Lagrangian immersions is a tangle invariant living in an appropriately defined Fukaya category of the pillowcase). Moreover, it will provide a mechanism for calculating the higher differentials in the spectral sequence, since they will be combinatorially computable via the Riemann mapping theorem for polygons. While much remains to be done to achieve this goal, we are given hope from the fact that we have already established the skein exact triangle for pillowcase homology in our work-in-progress.

Outline: We now summarize the results of the article. In Section 2 we recall and extend work of Abouzaid [1] to construct a Lagrangian-Floer theory for curves in surfaces, and outline the basic properties of the Maslov index.

In Section 3 we recall the construction of the pillowcase P as a quotient of \mathbb{R}^2 by $\mathbb{Z}^2 \ltimes \mathbb{Z}/2$. This is a 2-sphere with four singular points (*corners*). Motivated by the main result of our previous article [18], we fix a family $\{L_0^{\epsilon, g}\}_{\epsilon, g}$ of immersed circles with one double point in a certain regular homotopy class. We define a *restricted immersed 1-manifold in P* (roughly)

to be an immersion $L_1 : R \rightarrow P$, where either R is a circle and L_1 misses the corners of P , or R is an arc with endpoints mapping to the corners; see Definition 3.6. Choosing $L_0 = L_0^{\epsilon, g}$ transverse to a restricted lagrangian L_1 , we then define a chain complex $(C^\natural(L_0, L_1), \partial)$ with differential ∂ determined by immersed bigons in the smooth part of P with boundary lying on L_0 and L_1 , following Floer [16] and Abouzaid [1]. We show how to endow this complex with a relative $\mathbb{Z}/4$ grading, a variant of an idea due to Seidel [31].

In Section 4, we prove that the resulting Floer homology $HF(L_0, L_1)$ depends only on the homotopy classes of the restricted immersed curves L_0, L_1 (the result paraphrased as the theorem above). This result, together with some basic observations described in Section 5, provides a set of tools to calculate $HF(L_0, L_1)$.

In Section 6, we apply this construction to traceless character varieties of knots. We first recall that the traceless character variety of the pair $(S^2, \{a, b, c, d\})$ is the pillowcase P . The main theorem of [18] shows that if (X, K) is a knot in a 3-manifold and $(S^2, \{a, b, c, d\}) \subset (X, K)$ is a 2-sphere which separates (X, K) into a trivial 2-tangle in the 3-ball (D, U) and its complement (Y, T) , then generators of Kronheimer-Mrowka's reduced instanton knot complex can be identified with the intersection of $L_0^{\epsilon, g}$ and $L_1 : R(Y, T) \rightarrow P$. Hence, in favorable circumstances, to such a decomposition and an appropriate holonomy perturbation π we can assign the corresponding Floer homology $HF(L_0^{\epsilon, g}, L_1)$, which we denote by $H^\natural(Y, T, \pi)$. This leads us to make the Atiyah-Floer conjecture stated above (see Conjectures 6.3 and 6.5 for more precise statements).

We show in Section 7 that this conjecture holds for 2-bridge knots (where all differentials are zero in both complexes). Sections 8 and 9 establish some general properties of $R_\pi(Y, T)$, such as identifying the two boundary points, showing they are stable under holonomy perturbations, and map to the corners of the pillowcase. We also examine the effect of applying holonomy perturbations in a collar neighborhood of the separating 2-sphere; in particular, this is used to make L_0 and L_1 transverse.

In Sections 10 and 11 we turn to calculations for torus knots, which display a rich and complicated collection of examples. We find two appropriate perturbation curves in a useful tangle decomposition for any (p, q) torus knot, and prove (Theorem 10.1) that there exist perturbations π so that $R_\pi(Y, T)$ is a compact 1-manifold with two boundary points. We extend the work of [17], identifying $R_\pi(Y, T)$ and its image in the pillowcase, to give many calculations of $H^\natural(Y, T, \pi)$ for tangles associated to torus knots. We give examples where different tangle decompositions and perturbations of a knot yield the same Lagrangian-Floer homology, which agrees with reduced instanton homology. We give examples with non-zero differentials. The reader is encouraged to examine Figures 19 through 25 to get a feel for how calculations are carried out.

The upshot of our calculations is that the conjecture stated above holds for all the calculations of $H^\natural(Y, T)$ for which the corresponding instanton homology $I^\natural(S^3, K)$ is known, and is consistent with the conjectured equality of ranks of $I^\natural(S^3, K)$ and the Heegaard knot Floer homology of K when the instanton homology is unknown.

As an important final remark, we should say that while this article is rather lengthy we believe the results are quite natural and can be relatively easily understood through examples. Thus for the benefit of the reader we have included a running example which is illustrated in Figures 1, 5, 6, and 8, and with details of the resulting calculations in Section

3.9. Understanding this example, and its relation to the traceless character variety associated to a 2-tangle decomposition of the $(5, 11)$ torus knot (Section 11.1) should make our ideas quickly accessible.

Acknowledgements The first author gratefully acknowledges support from NSF grant DMS-0906258, NSF CAREER grant DMS-1150872, and an Alfred P. Sloan Research Fellowship. The third author gratefully acknowledges support from NSF grant DMS-1007196 and Simons foundation collaboration grant 278714.

The authors thank Juanita Pinzon-Caicedo and Yoshihiro Fukumoto for helpful discussions and assistance with writing the computer programs to produce the data and figures used in Section 11. They also thank Nikolai Saveliev, Matt Hogencamp for providing very useful insights, and Paul Seidel for pointing us to some influential references.

2. IMMERSED LAGRANGIAN FLOER THEORY ON A SURFACE

In [1], Abouzaid constructs a Lagrangian-Floer theory for *unobstructed immersed* curves in an oriented surface. In this section we recall his construction, adapted slightly for our purposes. We also recall and relate various versions of the Maslov index for curves and n -gons in a surface equipped with a line field.




FIGURE 1. An immersed circle L_1 in the 4-punctured 2-sphere.

2.1. Unobstructed immersed curves. Let S be a compact oriented surface, possibly with boundary, with infinite fundamental group.

Definition 2.1. An *unobstructed immersed arc* is an immersion $L : [0, 1] \rightarrow S$ which maps the endpoints to distinct points in the boundary of S , which is transverse to the boundary at its endpoints, and so that some (and hence every) lift of L to the universal cover of S is embedded.

An *unobstructed immersed circle* is an immersion of a circle $L : S^1 \rightarrow S$ so that each lift of the composite $\mathbb{R} \xrightarrow{e} S^1 \xrightarrow{L} S$ to the universal cover of S is a properly embedded line. Here $e(t) = \exp(2\pi it)$.

Either one of these is called an *unobstructed immersed curve*.

An immersion of a 1-manifold R to S is said to *contain a fishtail* if there is an interval $I \subset R$ whose endpoints are sent to the same point in S and so that the resulting loop is nullhomotopic. Lemma 2.2 of [1] shows that an immersed circle is unobstructed if and only if it is homotopically essential and contains no fishtails. Similarly, an immersed arc is unobstructed if and only if it contains no fishtails.

The reader can easily verify that the immersed circle illustrated in Figure 1 is essential and contains no fishtails, hence is unobstructed.

2.2. Intersection points. Let $L_0 : R_0 \rightarrow S$ and $L_1 : R_1 \rightarrow S$ be unobstructed immersed curves which intersect transversely.

Definition 2.2. Define an *intersection point* of L_0 and L_1 to be a pair $p = (r_0, r_1) \in R_0 \times R_1$ where $L_0(r_0) = L_1(r_1)$.

By transversality and compactness, there are only finitely many intersection points. We will frequently abuse notation and write “ $L_0 \cap L_1$ ” for the set of intersection points of L_0 with L_1 , or confuse $p = (r_0, r_1)$ with its image $L_0(r_0) = L_1(r_1)$ in S .

Define $C(L_0, L_1)$ to be the \mathbb{F}_2 vector space generated by the intersection points of L_0 and L_1 .

2.3. Line fields and the Maslov index. We recall some well known facts about the Maslov index in the 2-dimensional setting for the convenience of the reader, but also to make our conventions precise.

First, suppose that $\alpha : [0, 1] \rightarrow S^1$ is a smooth map. Define *the degree of α* to be the sum of the local degrees of α over the preimages of a regular value $e^{\delta i}$ for any small $\delta \geq 0$ chosen so that $\alpha(0), \alpha(1) \notin \{e^{t\delta i} \mid t \in (0, 1]\}$:

$$\deg(\alpha) := \sum_{x \in \alpha^{-1}(e^{\delta i})} \deg_x(\alpha)$$

($\deg_x(\alpha)$ denotes the sign of $d\alpha$ at x). Typically $\alpha(0) \neq 1 \neq \alpha(1)$ so that we may take $\delta = 0$. The integer $\deg(\alpha)$ has the property that it is additive under composition of paths, and invariant under homotopy of α rel endpoints. Define the degree of a continuous path to be the degree of any smooth approximation with the same endpoints.

Next, suppose that $\mathbf{P} \rightarrow B$ is a principal circle bundle over a space B and ℓ_0, ℓ are two sections. The section ℓ defines a trivialization $\mathbf{P} \cong_\ell B \times S^1$ sending ℓ to $1 \in S^1$. Given a path $\alpha : [0, 1] \rightarrow B$, define *the Maslov index of α with respect to ℓ_0 and ℓ* , denoted $\mu(\ell_0, \ell)_\alpha$, to be the degree of the composite

$$[0, 1] \xrightarrow{\alpha} B \xrightarrow{\ell_0} \mathbf{P} \cong_\ell \mathbf{B} \times \mathbf{S}^1 \xrightarrow{\pi} \mathbf{S}^1.$$

For the rest of this section we consider a pair (S, ℓ) , where S is an oriented Riemannian surface equipped with a line field ℓ (so S is either a torus or else S is not closed). Formally,

ℓ is a section of the projective tangent bundle $\mathbf{P}(T_*S)$, which we consider as a principal $S^1 = SO(2)/(\mathbb{Z}/2)$ bundle. Hence ℓ defines a trivialization $\mathbf{P}(T_*S) \cong_\ell S \times S^1$.

An immersion of a 1-manifold R in S defines a line field along R and hence a Maslov index for any path in R . More precisely, given an immersion $L_0 : R \rightarrow S$, the subspaces $dL_0(T_r R) \subset T_{L_0(r)} S$ of L define a section, which we denote ℓ_0 , of the pullback bundle $L_0^*(\mathbf{P}(T_*S)) \rightarrow R$. Taking account of the pullback of the line field ℓ , this yields the composite map

$$(1) \quad R \xrightarrow{\ell_0} L^*(\mathbf{P}(T_*S)) \cong_\ell R \times S^1 \xrightarrow{\pi} S^1$$

Definition 2.3. Given an immersion $L_0 : R \rightarrow S$ of a 1-manifold and a path $\alpha : [0, 1] \rightarrow R$, the *Maslov index* $\mu(L_0, \ell)_\alpha$, is defined to be $\mu(\ell_0, \ell)_\alpha$, the Maslov index of α with respect to ℓ_0 and ℓ . When clear from context this will be denoted $\mu(\alpha, \ell)$ or simply $\mu(\alpha)$.

As explicit examples (and to explain our conventions), if $S = \mathbb{R}^2$, ℓ is the horizontal line field, and R is the parabola $y = x^2$, then the path $\alpha : [-1, 1] \rightarrow R$ given by $t \mapsto (t, t^2)$ satisfies $\mu(\alpha) = 1$. If $\beta : [0, 1] \rightarrow R$ is given by $\beta(t) = (t, t^2)$ and $\gamma : [-1, 0] \rightarrow R$ is given by $\gamma(t) = (t, t^2)$, then $\mu(\beta) = 1$ and $\mu(\gamma) = 0$.

The basic properties of μ , including its dependence on the choice of the background line field ℓ on S , are well known and easily understood using obstruction theory. We summarize the facts we need in the following Proposition.

Proposition 2.4.

- i. If $L_0 : R \rightarrow S$ is an immersion of a 1-manifold and $\alpha, \beta : [0, 1] \rightarrow R$ are continuous paths with $\alpha(1) = \beta(0)$, then $\mu(L_0, \ell)_\alpha$ and $\mu(L_0, \ell)_\beta$ depend only on the homotopy classes of α and β rel boundary. Moreover, $\mu(L_0, \ell)_{\alpha*\beta} = \mu(L_0, \ell)_\alpha + \mu(L_0, \ell)_\beta$.
- ii. If $R = S^1$ and $\alpha(t) = e^{2\pi i t}$, $0 \leq t \leq 1$, then $\mu(L_0, \ell)_\alpha$ is unchanged by a regular homotopy of L_0 . More generally, if R is any 1-manifold and $\alpha : [0, 1] \rightarrow R$ arbitrary, then $\mu(L_0, \ell)_\alpha$ is unchanged by a regular homotopy of $L_0 : R \rightarrow S$ which leaves $\alpha(0)$ and $\alpha(1)$ and the tangent spaces of L_0 at these points stationary.
- iii. If ℓ' is any other line field, let $z \in [S, S^1] = H^1(S; \mathbb{Z}) = \text{Hom}(H_1(S), \mathbb{Z})$ denote the homotopy class of the difference map (i.e., $\ell'(s) = z(s)\ell(s)$). Here, we identify S^1 with \mathbb{RP}^1 , so that one rotation corresponds to a rotation of a line through an angle of π . If $\alpha : [0, 1] \rightarrow R$ is a loop, then $\mu(L_0, \ell')_\alpha = \mu(L_0, \ell)_\alpha + z(L_0 \circ \alpha)$.

We next define the triple index $\tau(\ell_0, \ell_1, \ell)$.

Definition 2.5. Suppose that $s \in S$. Given a pair ℓ_0, ℓ_1 of transverse 1-dimensional subspaces of $T_s S$, let $\ell_t, t \in [0, 1]$ be the *shortest clockwise path* from ℓ_0 to ℓ_1 . Then define the *triple index*

$$\tau(\ell_0, \ell_1, \ell) = -\mu(\ell_t, \ell)_{[0,1]} \in \{0, 1\}$$

When ℓ_0, ℓ_1 and ℓ are pairwise transverse, $\tau(\ell_0, \ell_1, \ell)$ is equal to 1 if ℓ_0 passes through ℓ when rotating ℓ_0 negatively (clockwise) to ℓ_1 ; otherwise $\tau(\ell_0, \ell_1, \ell) = 0$. See Figure 2.

The triple index has the following properties: if ℓ_0, ℓ_1, ℓ_2 are pairwise transverse, then

$$(2) \quad \tau(\ell_1, \ell_2, \ell_0) = \tau(\ell_0, \ell_1, \ell_2) \text{ and } \tau(\ell_1, \ell_0, \ell) = 1 - \tau(\ell_0, \ell_1, \ell_2)$$




FIGURE 2.

Let $L_k : R_k \rightarrow S$, $k = 0, 1, 2, \dots, n-1$, be a sequence of pairwise transverse unobstructed immersed curves. Let p_k be an intersection point of L_{k-1} and L_k for $k = 1, \dots, n$. We consider the indices cyclically ordered, so that $L_n = L_0$ and $p_0 = p_n$. Figure 3 illustrates the notation.




FIGURE 3. A 5-gon

Definition 2.6. Given an ordered list L_0, L_1, \dots, L_{n-1} of pairwise transverse unobstructed immersed curves and a sequence of intersection points $p_n = p_0, p_1, \dots, p_{n-1}$ with $p_i \in L_{i-1} \cap L_i$, define $\pi_1(p_1, \dots, p_n)$ to be the homotopy rel boundary classes of n -tuples of paths $(\gamma_0, \dots, \gamma_{n-1})$, where γ_k is a path in R_k from (the first coordinate of the) intersection point p_k to p_{k+1} .

The reader will notice that $p_n = p_0$ is placed last in the notation, in contrast to the order of the L_i (and γ_i) where L_0 is placed first. This is motivated by the fact that it is often useful to treat $p_0 = p_n$ differently than the rest of the p_k , for example in the definition of the differential and, more generally, the A_∞ structure on the Fukaya category of S (see Definition 2.13 below). In particular, this ensures that notation to be introduced below for an n -gon from (p_1, \dots, p_{n-1}) to $p_n = p_0$ is as simple as possible.

Definition 2.7. Define $\pi_2(p_1, \dots, p_n)$ to be homotopy classes of pairs $(u, (\gamma_0, \dots, \gamma_{n-1}))$ where $(\gamma_0, \dots, \gamma_{n-1}) \in \pi_1(p_1, \dots, p_n)$, and $u : D^2 \rightarrow S$ is a continuous map so that the (counterclockwise) boundary of D^2 is sent to the loop

$$\left((L_0 \circ \gamma_0) * \dots * (L_{n-1} \circ \gamma_{n-1}) \right)^{-1}.$$

There is a well defined forgetful map $\pi_2(p_1, \dots, p_n) \rightarrow \pi_1(p_1, \dots, p_n)$ obtained by sending the equivalence class of $(u, (\gamma_0, \dots, \gamma_{n-1}))$ to that of $(\gamma_0, \dots, \gamma_{n-1})$.

There is a well defined function

$$\theta : \pi_1(p_1, \dots, p_n) \rightarrow H_1(S), \quad \theta(\gamma_0, \dots, \gamma_{n-1}) = -[(L_0 \circ \gamma_0) * \dots * (L_{n-1} \circ \gamma_{n-1})]$$

which vanishes on the image of the forgetful map $\pi_2(p_1, \dots, p_n) \rightarrow \pi_1(p_1, \dots, p_n)$.

We use a “label clockwise, but orient the boundary of a disk counterclockwise” convention. This is why the inverse of the loop $L_0(\gamma_0) * \dots * L_{n-1}(\gamma_{n-1})$ appears in Definition 2.7, and that the negative sign is used in the definition of θ .

Given an intersection point p_k of L_{k-1} and L_k , $\pi_2(p_k, p_k)$ is a group. If $\pi_2(p_1, \dots, p_n)$ is non-empty, then $\pi_2(p_k, p_k)$ acts on $\pi_2(p_1, \dots, p_n)$ by attaching $\tau \in \pi_2(p_k, p_k)$ to $\phi \in \pi_2(p_1, \dots, p_n)$ along the vertex p_k to form a new map of the disk $\tau \cdot \phi \in \pi_2(p_1, \dots, p_n)$.

In the case of a pair (L_0, L_1) , it is easy to check that $\pi_2(p_1, p_1)$ acts freely and transitively on $\pi_2(p_1, p_2)$ for any p_2 for which $\pi_2(p_1, p_2)$ is non-empty.

Definition 2.8. Given an ordered list L_0, L_1, \dots, L_{n-1} of pairwise transverse unobstructed immersed curves, intersection points $p_k \in L_{k-1} \cap L_k$, and an n -tuple of paths $(\gamma_0, \dots, \gamma_{n-1})$ representing a class in $\pi_1(p_1, \dots, p_n)$, define

$$\text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = 1 - \sum_{k=0}^{n-1} (\mu(L_k, \ell)_{\gamma_k} + \tau(L_k, L_{k-1}, \ell)_{p_k})$$

We describe a few equivalent formulas for Mas_ℓ . First notice that letting α_k denote the reverse of the path γ_k ,

$$\alpha_k(t) = \gamma_k(1-t),$$

then $\mu(L_k, \ell)_{\gamma_k} = -\mu(L_k, \ell)_{\alpha_k}$. Next, $-\tau(L_k, L_{k-1}, \ell)_{p_k}$ is equal to $\mu(M_k(t), \ell)_{[0,1]}$, where $M_k(t)$ is the path of lines in $T_{p_k}S$ obtained as the shortest clockwise rotation of $T_{p_k}L_k$ to $T_{p_k}L_{k-1}$. Hence

$$(3) \quad \text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = 1 + \sum_{k=0}^{n-1} (\mu(L_k, \ell)_{\alpha_k} + \mu(M_k(t), \ell)_{[0,1]})$$

which, by path additivity of the Maslov index, is one greater than the Maslov index with respect to the line field ℓ of the continuous loop in the projective tangent bundle of S :

$$\mathbb{M} := TL_{n-1}|_{\alpha_{n-1}} * M_{n-1} * TL_{n-2}|_{\alpha_{n-2}} * M_{n-2} * \dots * TL_0|_{\alpha_0} * M_0,$$

i.e.,

$$\text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = 1 + \mu(\mathbb{M}, \ell).$$

As mentioned before, p_0 plays a special role, which motivates taking the continuous loop obtained by rotating clockwise at $p_k, k > 0$, but rotating *counterclockwise* at p_0 :

$$\mathbb{A} := TL_{n-1}|_{\alpha_{n-1}} * M_{n-1} * TL_{n-2}|_{\alpha_{n-2}} * M_{n-2} * \dots * TL_0|_{\alpha_0} * N_0$$

where $N_0(t)$ is the shortest counterclockwise rotation of $T_{p_0}L_0$ to $T_{p_0}L_{n-1}$ in $T_{p_0}S$. Then using (2),

$$(4) \quad \text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = \mu(\mathbb{A}, \ell),$$

justifying the use of the notation Mas_ℓ .

Proposition 2.9. *Given an ordered set L_0, \dots, L_{n-1} of pairwise transverse unobstructed immersed curves and intersection points p_k of L_{k-1} and L_k , then Mas_ℓ defines a function $\pi_1(p_1, \dots, p_n) \rightarrow \mathbb{Z}$. If ℓ' is another line field and $z \in H^1(S) = [S, S^1]$ is the difference class, then*

$$\text{Mas}_{\ell'}(\gamma_0, \dots, \gamma_{n-1}) - \text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = z(\theta(\gamma_0, \dots, \gamma_{n-1})).$$

In particular, Mas_ℓ depends only on the homotopy class of ℓ , and the composite

$$\pi_2(p_1, \dots, p_n) \rightarrow \pi_1(p_1, \dots, p_n) \xrightarrow{\text{Mas}_\ell} \mathbb{Z}$$

is independent of the choice of line field ℓ .

Furthermore, Mas_ℓ has the properties:

- i. (Splicing) *If q is another intersection point of L_0 and L_k , and $\gamma'_0, \gamma''_0, \gamma'_k, \gamma''_k$ are paths so that $\gamma'_0(1) = q = \gamma''_0(0)$, $\gamma'_k(1) = q = \gamma''_k(0)$, $\gamma_0 = \gamma'_0 * \gamma''_0$, and $\gamma_k = \gamma'_k * \gamma''_k$, then*

$$\text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = \text{Mas}_\ell(\gamma''_0, \gamma_1, \dots, \gamma_{k-1}, \gamma'_k) + \text{Mas}_\ell(\gamma''_k, \gamma_{k+1}, \dots, \gamma_{n-1}, \gamma'_0).$$

- ii. (Path reversal) *Let $\alpha_k(t) = \gamma_k(1 - t)$. Then*

$$\text{Mas}_\ell(\alpha_{n-1}, \alpha_{n-2}, \dots, \alpha_0) = 2 - n - \text{Mas}_\ell(\gamma_0, \gamma_1, \dots, \gamma_{n-1}).$$

- iii. (Cyclic invariance) $\text{Mas}_\ell(\gamma_0, \gamma_1, \dots, \gamma_{n-1}) = \text{Mas}_\ell(\gamma_1, \dots, \gamma_{n-1}, \gamma_0)$.

Proof. The homotopy invariance property of the Maslov index μ and Equation (4) shows that $\text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1})$ depends only on the homotopy class of ℓ and the class of $(\gamma_0, \dots, \gamma_{n-1})$ in $\pi_1(p_1, \dots, p_n)$.

If ℓ, ℓ' are two different line fields and $z \in H^1(S)$ is their difference class, Proposition 2.4 gives

$$\text{Mas}_{\ell'}(\gamma_0, \dots, \gamma_{n-1}) - \text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1}) = \mu(\mathbb{A}, \ell') - \mu(\mathbb{A}, \ell) = z(\theta(\gamma_0, \dots, \gamma_{n-1})).$$

If $(\gamma_0, \dots, \gamma_{n-1})$ lies in the image of $\pi_2(p_1, \dots, p_n) \rightarrow \pi_1(p_1, \dots, p_n)$, then $\theta(\gamma_0, \dots, \gamma_{n-1}) = 0$ in $H_1(S)$, and hence $\text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1})$ is independent of ℓ .

The three properties are easily checked using path additivity of $\mu(-, \ell)$ and the identity $\tau(\ell_0, \ell_1, \ell) + \tau(\ell_1, \ell_0, \ell) = 1$. \square

2.4. Immersed polygons. Define an n -gon in \mathbb{R}^2 to be a pair $(D, (\beta_0, \beta_1, \dots, \beta_{n-1}))$ where $D \subset \mathbb{R}^2$ is a closed topological disc, and $(\beta_0, \beta_1, \dots, \beta_{n-1})$ is a sequence of smooth embeddings of the unit interval in \mathbb{R}^2 with image in the boundary of D so that

- i. $\beta_k(1) = \beta_{k+1}(0)$ for $k = 0, \dots, n$ (with $\beta_n = \beta_0$),
- ii. the composite path $\beta_0 * \dots * \beta_{n-1}$ forms an embedded simple closed curve in \mathbb{R}^2 , which forms the *clockwise* boundary of D ,
- iii. the β_k meet transversely at their endpoints.

Definition 2.10. Given a pair $(S, (L_0, \dots, L_{n-1}))$, where S is an oriented surface and $L_k : R_k \rightarrow S$, $k = 0, \dots, n-1$ are pairwise transverse immersions of 1-manifolds into S , define an *immersed n -gon in S for the ordered n -tuple (L_0, \dots, L_{n-1}) through the points (p_1, \dots, p_n)* to be a triple consisting of

- i. an n -gon in \mathbb{R}^2 , $(D, (\beta_0, \beta_1, \dots, \beta_{n-1}))$,
- ii. a representative n -tuple of paths $(\gamma_0, \dots, \gamma_{n-1})$ for a class in $\pi_1(p_1, \dots, p_n)$,
- iii. an orientation preserving immersion $u : D \rightarrow S$ satisfying $u \circ \beta_k = L_k \circ \gamma_k$.

We use the brief notation u for the triple

$$((D, (\beta_0, \dots, \beta_{n-1})), (\gamma_0, \dots, \gamma_{n-1}), u : D \rightarrow S),$$

and we call this an *immersed n -gon for the ordered n -tuple (L_0, \dots, L_{n-1}) from (p_1, \dots, p_{n-1}) to $p_0 = p_n$* .

Define the *Maslov index of an immersed n -gon in S* by

$$\text{Mas}(u) = \text{Mas}_\ell(\gamma_0, \dots, \gamma_{n-1})$$

for any line field ℓ . Since an immersed n -gon represents an element of $\pi_2(p_1, \dots, p_n)$, it follows from Proposition 2.9 that $\text{Mas}(u)$ is independent of the choice of line field ℓ .

Figure 4 indicates the corresponding model examples of 2-gons and 3-gons in $S = \mathbb{R}^2$:




FIGURE 4. A 2-gon of Maslov index 1 from p to q for the ordered pair (L_0, L_1) and a 3-gon of Maslov index 0 from (p, q) to r for the ordered triple (L_0, L_1, L_2) .

Notice that an n -gon in \mathbb{R}^2 is just a special (embedded) case of an immersed n -gon in the surface $S = \mathbb{R}^2$. As such, its Maslov index can be easily computed by taking, for example, any line field of constant slope. For example, the 5-gon of Figure 3 has Maslov index -1 . One can see this by taking ℓ to be the vertical line field. Then $\mu(L_k, \ell)_{\gamma_k} = 0$ for $i = 0, 1, 2, 3$ and $\mu(L_4, \ell)_{\gamma_4} = -2$. Also $\tau(L_k, L_{k-1}, \ell)_{p_k}$ equals 1 for $k = 0, 2, 3, 4$ and $\tau(L_1, L_0, \ell)_{p_1} = 0$, so $\text{Mas} = 1 - (-2 + 4) = -1$. This calculation easily generalizes to yield the following proposition.

Proposition 2.11. *Let $(D, \beta_0, \dots, \beta_{n-1})$ be an n -gon in \mathbb{R}^2 . Let $\kappa(D) \in \{0, \dots, n\}$ denote the number of non-convex corners of D .*

Then any immersed n -gon $u : D \rightarrow S$ satisfies

$$\text{Mas}(u) = 3 - n + \kappa(D).$$

□

2.5. The differential and A_n maps. We recall the definitions of the differential and maps μ_n next. Our goal is to relate a Lagrangian-Floer theory to singular instanton homology, which is the homology of a chain complex rather than a cochain complex, and hence our orientation conventions differ slightly from the similar constructions in the literature which typically produce cochain complexes.

Definition 2.12. Fix an ordered n -tuple $(L_0, L_1, \dots, L_{n-1})$ of distinct unobstructed pairwise transverse immersed curves in S and intersection points p_k of L_{k-1} and L_k .

Two immersed n -gons in S through (p_1, \dots, p_n) , $u : D \rightarrow S, u' : D' \rightarrow S$ are called *equivalent* if there is an orientation preserving diffeomorphism $\psi : D \rightarrow D'$ so that $u = u' \circ \psi$.

The set of equivalence classes of immersed n -gons with Maslov index $3 - n$ for the ordered n -tuple (L_0, \dots, L_{n-1}) through (p_1, \dots, p_n) is denoted by $\mathcal{M}_{L_0, \dots, L_{n-1}}(p_1, \dots, p_n)$, or simply by $\mathcal{M}(p_1, \dots, p_n)$ when the order of the L_k is clear from context.

When $n \geq 3$, the list (p_1, \dots, p_n) determines the order (L_0, \dots, L_{n-1}) . Cyclically permuting the n -tuple (L_0, \dots, L_{n-1}) and the points (p_1, \dots, p_n) preserves immersed bigons and the Maslov index, and hence

$$\mathcal{M}_{L_0, L_1, \dots, L_{n-1}}(p_1, \dots, p_n) = \mathcal{M}_{L_1, \dots, L_{n-1}, L_0}(p_2, \dots, p_n, p_1).$$

When $n = 2$, care must be taken with the ordering since the ordered pair (p_0, p_1) does not determine the order of L_0, L_1 . In particular, $\mathcal{M}_{L_0, L_1}(p, q) = \mathcal{M}_{L_1, L_0}(q, p)$, but these are different from $\mathcal{M}_{L_1, L_0}(p, q) = \mathcal{M}_{L_0, L_1}(q, p)$.

Given a finite set X , let $\#X \in \mathbb{F}_2$ denote the number of elements of X mod 2. Recall that $C(L_0, L_1)$ is defined to be the free \mathbb{F}_2 vector space on the intersection points of L_0 and L_1 .

Definition 2.13. Fix an ordered n -tuple (L_0, \dots, L_{n-1}) of unobstructed pairwise transverse curves in S . Suppose $n - 1$ intersection points p_k of L_{k-1} and L_k are given for $k = 1, \dots, n - 1$, with the property that for every intersection point q of L_0 and L_{n-1} , $\mathcal{M}_{L_0, \dots, L_{n-1}}(p_1, \dots, p_{n-1}, q)$ is finite.

Define

$$\mu_{n-1}(p_1, \dots, p_{n-1}) = \sum_{q \in L_0 \cap L_{n-1}} (\# \mathcal{M}_{L_0, \dots, L_{n-1}}(p_1, \dots, p_{n-1}, q)) q \text{ in } C(L_0, L_{n-1}).$$

If $\mathcal{M}_{L_0, \dots, L_{n-1}}(p_1, \dots, p_n)$ is finite for all choices of intersection points (p_1, \dots, p_{n-1}) and p_n , then μ_{n-1} defines a linear map:

$$\mu_{n-1} : C(L_0, L_1) \otimes C(L_1, L_2) \otimes \dots \otimes C(L_{n-2}, L_{n-1}) \rightarrow C(L_0, L_{n-1}).$$


Most important for us is the map μ_1 , which we also denote by ∂ . Explicitly

$$(5) \quad \partial : C(L_0, L_1) \rightarrow C(L_0, L_1), \quad \partial p = \sum_{q \in L_0 \cap L_1} \# \mathcal{M}_{L_0, L_1}(p, q) q$$

Recall that we call representatives of $\mathcal{M}_{L_0, L_1}(p, q)$ 2-gons from p to q for (L_0, L_1) , so ∂p is the linear combination of intersection points q weighted by the mod 2 count of 2-gons with Maslov index 1 from p to q .

Since they occur frequently, we call a 2-gon with Maslov index 1 a *bigon*. Note that a bigon from p to q for the ordered pair (L_0, L_1) is also a bigon from q to p for (L_1, L_0) , but is not a bigon from q to p for (L_0, L_1) . See the paragraph following Definition 2.12.

Figure 5 shows the curve L_1 of Figure 1 and another unobstructed curve L_0 . These intersect in eight points. Two intersection points p, q are indicated, and the image of a bigon from p to q for the pair (L_0, L_1) is shaded. The reader should check that there is precisely one other bigon between L_0 and L_1 , joining a different pair of intersection points.

FIGURE 5. A bigon from p to q .

The following is proved in Abouzaid's article [1]. In that article, coefficients are taken in a Novikov ring (over \mathbb{Z}) to account for the possibility that $\mathcal{M}_{L_0, L_1}(q, p)$ is infinite. Since this will not be the case in our applications, we set the Novikov variable t equal to 1 and reduce the coefficients from \mathbb{Z} to \mathbb{F}_2 .

Theorem 2.14 (Abouzaid [1]). *Let (L_0, L_1) be a pair of unobstructed transverse immersed curves in S , and assume that $\mathcal{M}_{L_0, L_1}(p, q)$ is finite for all intersection points p, q of L_0 and L_1 . Then $\partial : C(L_0, L_1) \rightarrow C(L_0, L_1)$ satisfies $\partial^2 = 0$. \square*

For example, the chain complex $C(L_0, L_1)$ for the pair (L_0, L_1) illustrated in Figure 5 (see also Figures 6 and 8) is generated by the eight intersection points of L_0 and L_1 . Two bigons (one of which is shaded in Figure 5) define a non-trivial differential ∂ of rank 2. The resulting homology has rank four.

More generally, Abouzaid proves in [1] that the μ_n , $n \geq 2$ satisfy the A_n relations for all n when (L_0, \dots, L_{n-1}) are pairwise transverse unobstructed immersed curves with no triple points.

We will only use the A_2 and A_3 relations in the present article and so we write them out explicitly. (We refer to [1, 4] for the formulas for the A_n relations.) The A_2 relation says

$$(6) \quad \mu_2(\mu_1(x), y) + \mu_2(x, \mu_1(y)) + \mu_1(\mu_2(x, y)) = 0,$$

and the A_3 relation says

$$(7) \quad \begin{aligned} &\mu_3(\mu_1(x), y, z) + \mu_3(x, \mu_1(y), z) + \mu_3(x, y, \mu_1(z)) + \mu_2(\mu_2(x, y), z) \\ &\quad + \mu_2(x, \mu_2(y, z)) + \mu_1(\mu_3(x, y, z)) = 0 \end{aligned}$$

These hold when all the sets of equivalence classes of n -gons of Maslov index $3 - n$, $n = 2, 3, 4$ which appear in the formulas defining each term in Equation (6) or (7) are finite.

3. RESTRICTED LAGRANGIANS IN THE PILLOWCASE

In this section, we will apply the constructions of Section 2 to the pillowcase.

3.1. The pillowcase. The *pillowcase* P is the quotient of the torus by the hyperelliptic involution. It is a topological 2-sphere with four singular points corresponding to the four fixed points of the involution. For concreteness, define P to be the quotient of \mathbb{R}^2 by the group of orientation preserving isometries generated by the maps

$$(\gamma, \theta) \mapsto (\gamma + 2\pi, \theta), \quad (\gamma, \theta) \mapsto (\gamma, \theta + 2\pi), \quad (\gamma, \theta) \mapsto (-\gamma, -\theta)$$

(this group is a semi-direct product $\mathbb{Z}^2 \ltimes \mathbb{Z}/2$). The quotient map is a branched covering

$$(8) \quad \mathbb{R}^2 \rightarrow P.$$

A fundamental domain for the action is given by the rectangle $(\gamma, \theta) \in [0, \pi] \times [0, 2\pi]$. We will frequently specify a point in P by giving its coordinates $(\gamma, \theta) \in \mathbb{R}^2$. We refer to points in $(\pi\mathbb{Z})^2$ as *lattice points*. The four singular points of P , which we call the corners, make up the image of the lattice points. Our theory will take place in the complement of the corners, so it is convenient to adopt the notation $P^* = (\mathbb{R}^2 \setminus (\pi\mathbb{Z})^2) / (\mathbb{Z}^2 \ltimes \mathbb{Z}/2)$. Note that P^* inherits an orientation and a symplectic structure from the standard orientation and symplectic structure $d\gamma \wedge d\theta$ on $\mathbb{R}^2 \setminus (\pi\mathbb{Z})^2$ via the branched covering (8).

The pillowcase P is illustrated in two ways in Figure 6. One should view the figure on the left as obtained by folding the fundamental domain $[0, \pi] \times [0, 2\pi]$ for the branched cover (8), illustrated on the right, along $[0, \pi] \times \{\pi\}$ and making identifications along the edges as indicated. The front face is the image of $[0, \pi] \times [0, \pi]$ and the back face is the image of $[0, \pi] \times [\pi, 2\pi]$, upside down. In Figure 6 we have also indicated the immersed circle L_1 of Figure 1.




FIGURE 6. Two depictions of the pillowcase P . The immersed circle L_1 of Figure 1 is indicated on the right.

3.2. A line field on the pillowcase. To apply the Maslov index constructions described in Section 2, a line field on P^* is needed. We fix a line field ℓ_{inst} in a particular homotopy class, so that the $\mathbb{Z}/4$ grading we construct below using the Maslov index matches the $\mathbb{Z}/4$

grading on singular instanton knot homology. The connection to gauge theory is explained in Section 6.

The line field ℓ_{inst} is somewhat complicated to depict or calculate with, as it twists along the edges of the pillowcase. Our approach in calculations is to keep track of the mod 4 Maslov index information determined by ℓ_{inst} by instead using a pair (ℓ, z) , where ℓ is a simple (constant slope) line field and $z \in H^1(P^*; \mathbb{Z}/4)$ keeps track of the extra twisting of ℓ_{inst} relative to ℓ , as described in Proposition 2.4.

Any constant slope line field on \mathbb{R}^2 is invariant under the $\mathbb{Z}^2 \ltimes \mathbb{Z}/2$ action and hence its restriction to $\mathbb{R}^2 \setminus (\pi\mathbb{Z})^2$ descends to a line field on P^* . Call such a line field a *constant slope* line field on P^* . We will make frequent use of the slope one line field on \mathbb{R}^2 , and hence we give it the label ℓ_1 .

Definition 3.1. Let $z \in H^1(P^*; \mathbb{Z}/4) = \text{Hom}(H_1(P^*), \mathbb{Z}/4)$ denote the unique cohomology class which assigns $1 \in \mathbb{Z}/4$ to each small loop circling a corner counterclockwise.

Immersed circles $\gamma : S^1 \rightarrow P^*$ satisfy $\mu(\gamma, \ell_1) \equiv z(\gamma) \pmod{2}$. They need not be equal modulo 4, however. For example, if γ is the boundary of a smoothly embedded disk in P^* , then $\mu(\gamma, \ell_1) = \pm 2$ and $z(\gamma) = 0$. For a small embedded loop encircling one corner of P counterclockwise, $\mu(\gamma, \ell_1) = 1 = z(\gamma)$. For the curve L_1 illustrated in Figure 1 and the right in Figure 6, $\mu(L_1, \ell_1) = 0 = z(L_1)$. For the curves $L_0^{\epsilon, g}$ depicted in Figure 7, $\mu(L_0^{\epsilon, g}, \ell_1) = 0 = z(L_0^{\epsilon, g})$.

Definition 3.2. Fix a map $\tilde{z} : P^* \rightarrow S^1$ so that its class $\tilde{z} \in H^1(P^*; \mathbb{Z}) = [P^*, S^1]$ is a lift of the class z . Think of S^1 as acting freely and transitively on lines in \mathbb{R}^2 . Define the *instanton line field* by:

$$(9) \quad \ell_{\text{inst}}(p) = \tilde{z}(p)\ell_1(p).$$

The homotopy class of the line field ℓ_{inst} depends on choice of lift \tilde{z} , as do Maslov indices computed using ℓ_{inst} , however,

$$(10) \quad \mu(L, \ell_{\text{inst}}) \equiv \mu(L, \ell_1) + z(L) \pmod{4}.$$

3.3. Proper arcs in P .

Definition 3.3. Define a *proper immersion* of an interval $L : I \rightarrow P$ to be the image under the branched cover (8) of a smooth immersion $\tilde{L} : I \rightarrow \mathbb{R}^2$ which takes the two endpoints of the interval to $(\pi\mathbb{Z})^2$ and the interior to $\mathbb{R}^2 \setminus (\pi\mathbb{Z})^2$. We call the slopes of \tilde{L} at the endpoints (which are determined by L) the *limiting slopes of L* . Note that a proper immersion cannot spiral infinitely many times as it limits to a corner.

In order to easily apply the results of [1], it is convenient to work in a compact surface with boundary. It will suffice for our purposes to simply remove a small neighborhood of the corners. More precisely, given some small $\delta > 0$, let $\bar{P}_\delta \subset P$ denote the image under the branched cover (8) of the subspace of \mathbb{R}^2 obtained by removing open δ neighborhoods of the lattice points.

If R is a compact 1-manifold with boundary and $L_1 : R \rightarrow P$ is a proper immersion (as defined above) on each arc, and maps each circle of R into P^* , then for $\delta > 0$ small enough so that the δ disks miss the circle components, $L_1(R) \cap \bar{P}_\delta$ is a properly immersed compact

1-manifold in the compact surface \overline{P}_δ . In the following, we will typically write \overline{P} instead of \overline{P}_δ , with the understanding that δ is chosen small enough to miss circle components and result in arc components transverse to the boundary.

3.4. Perturbation functions and a family of isotopies of P . Let

$$\mathcal{X} = \{f \in C^\infty(\mathbb{R}, \mathbb{R}) \mid f(x + 2\pi) = f(x), f \text{ odd}\}.$$

We call this the *space of perturbation functions*. It is a vector space, and is preserved by pre-composition by $x \mapsto x + \pi$. In particular, $f(\pi) = 0$ for all $f \in \mathcal{X}$. The sine function is a member of \mathcal{X} .

The usual terminology in the literature describes perturbation data as a choice of an embedded solid torus and a conjugation invariant function on $SU(2)$, which together are used to define a gauge invariant perturbation of the Chern-Simons functional. In our notation, an element $f \in \mathcal{X}$ is the derivative of such a conjugation invariant function on $SU(2)$, restricted to the maximal torus. The function f determines the effect on the critical set of Chern-Simons function. More precisely, f determines which flat connections on the complement of the perturbation solid torus extend to be perturbed flat on the solid torus, so it is more convenient for us to refer to these functions in our perturbation data. See Section 8.2 for more details.

We associate, to each perturbation function $g \in \mathcal{X}$, an isotopy of the pillowcase by:

$$(11) \quad c_g : P \times I \rightarrow P, \quad c_g((\gamma, \theta), s) = (\gamma, \theta + sg(\gamma)).$$

Since $c_{-g}(c_g(p, s), s) = p$, $c_g(-, s)$ is a homeomorphism and hence c_g is an isotopy starting at the identity. Notice that c_g fixes the left and right edges of the pillowcase.

The formula (11) shows that c_g lifts to a Hamiltonian isotopy of \mathbb{R}^2 which is $\mathbb{Z} \ltimes \mathbb{Z}/2$ invariant and fixes the vertical lines $\{x = n\pi\}$. In particular, we can think of c_g as a Hamiltonian isotopy of P^* , or of the orbifold P .

3.5. The family $L_0^{\epsilon, g}$ of immersed circles in the pillowcase. In the applications to singular instanton homology in Section 6, we show that a 2-tangle decomposition of a knot gives rise to two unobstructed immersed curves L_0, L_1 in P^* , which in turn define a chain complex $C(L_0, L_1)$ as in Section 2. Identification of the immersed circle L_0 , which depends on a parameter $\epsilon \neq 0$, was accomplished in [18, Theorem 7.1]. In order to ensure we can choose L_0 transverse to L_1 we enlarge the family of L_0 to include the isotopies described above.

Let $\Delta \subset P$ denote the arc of slope one, i.e., the diagonal arc

$$(12) \quad \Delta = \{(\gamma, \gamma) \mid \gamma \in [0, \pi]\}.$$

Definition 3.4. Fix an $\epsilon > 0$ and $g \in \mathcal{X}$. Let $L_0^{\epsilon, g} : S^1 \rightarrow P^*$ denote the immersion given as the composite of the map

$$(13) \quad \tilde{L}_0^{\epsilon, g}(t) = (t + \epsilon \sin(t) + \frac{\pi}{2}, t - \epsilon \sin(t) + \frac{\pi}{2} + g(t + \epsilon \sin(t) + \frac{\pi}{2})), \quad t \in [0, 2\pi]$$

with the branched cover $\mathbb{R}^2 \rightarrow P$ of Equation (8), so

$$(14) \quad L_0^{\epsilon, g} : [0, 2\pi] \xrightarrow{\tilde{L}_0^{\epsilon, g}} \mathbb{R}^2 \rightarrow P^*.$$

The image of $L_0^{\epsilon,g}$ in P^* for $g = 0$ and ϵ small is illustrated in Figure 7 and also in Figure 5. As ϵ and g approach zero, $L_0^{\epsilon,g}$ limits to a generically 2-1 map onto Δ , with two points mapping to corners.




FIGURE 7. The curve $L_0^{\epsilon,0}$ in P .

Note that $L_0^{\epsilon,g}(t) = c_g(L_0^{\epsilon,0}(t), 1)$, so that $L_0^{\epsilon,g}$ is isotopic to $L_0^{\epsilon,0}$. In particular, the family of immersions $L_0^{\epsilon,g}$ for $\epsilon > 0$ small are self-transverse with one double point. Furthermore, $L_0^{\epsilon,g}$ is an unobstructed circle in the sense of Definition 2.1.

The following easily proved genericity lemma says that given any unobstructed immersed curve L_1 , arbitrarily small ϵ, g can be found so that L_1 and $L_0^{\epsilon,g}$ are transverse.

Lemma 3.5. *Given an unobstructed immersed circle or arc L_1 , there exist $\epsilon > 0$ and $\delta > 0$ arbitrarily close to zero so that, with $g(x) = \delta \sin(x)$, $L_0^{\epsilon,g}$ and L_1 are transverse. \square*

3.6. Restricted immersed arcs and circles. To complete the construction of a $\mathbb{Z}/4$ relatively graded chain complex we refine the notion of an unobstructed curve. Experts will recognize this notion as a $\mathbb{Z}/4$ variant of Seidel's notion of a *graded Lagrangian* ([31, 4]).

Definition 3.6.

A *restricted immersed circle in P^** is an unobstructed immersed circle $L_1 : S^1 \rightarrow P^*$ which satisfies $\mu(L_1(S^1), \ell_{\text{inst}}) \equiv 0 \pmod{4}$, or, equivalently, $\mu(L_1(S^1), \ell_1) + z(L_1(S^1)) \equiv 0 \pmod{4}$.

A *restricted immersed arc in P* is a proper immersion on an interval (in the sense of Definition 3.3) $L_1 : I \rightarrow P$ such that $L_1(I) \cap \overline{P}_\delta$ is unobstructed for small $\delta > 0$.

A *restricted immersed curve* is either a restricted immersed circle or a restricted immersed arc.

The curves $L_0^{\epsilon,g}$ and the curve L_1 of Figure 1 are restricted immersed circles. An embedded circle L encircling one corner of P counterclockwise is unobstructed but not restricted since $\mu(L, \ell_1) + z(L) = 2$. The image of any straight line segment in \mathbb{R}^2 joining two lattice points whose interior misses the lattice is mapped via the branched cover (8) to a restricted immersed arc.

3.7. A relative $\mathbb{Z}/4$ grading. We revisit the notation and constructions of Section 2 in the context of restricted immersed curves. Recall that for simplicity we write $L_k \cap L_j$ for the set of intersection points of L_k with L_j (see Definition 2.2).

Definition 3.7. Given an ordered list (L_0, \dots, L_{n-1}) of pairwise transverse restricted immersed curves, define

$$gr_{L_0, L_1, \dots, L_{n-1}} : (L_0 \cap L_1) \times (L_1 \cap L_2) \times \dots \times (L_{n-2} \cap L_{n-1}) \times (L_{n-1} \cap L_0) \rightarrow \mathbb{Z}/4$$

by

$$gr_{L_0, L_1, \dots, L_{n-1}}(p_1, \dots, p_n) = \text{Mas}_{\ell_{\text{inst}}}(\gamma_0, \dots, \gamma_{n-1}) \pmod{4}$$

for any choice of $(\gamma_0, \dots, \gamma_{n-1}) \in \pi_1(p_1, \dots, p_n)$.

Proposition 3.8. *Given an ordered list (L_0, \dots, L_{n-1}) of pairwise transverse restricted immersed curves,*

- i. *$gr_{L_0, L_1, \dots, L_{n-1}}(p_1, \dots, p_n)$ is independent of the choice of $(\gamma_0, \dots, \gamma_n)$ and is invariant under simultaneous cyclic permutations of L_0, L_1, \dots, L_{n-1} and $p_1, \dots, p_{n-1}, p_n = p_0$.*
- ii. *If q is another intersection point of L_0 and L_k , then*

$$\begin{aligned} &gr_{L_0, L_1, \dots, L_{n-1}}(p_1, \dots, p_n) \\ &= gr_{L_0, L_1, \dots, L_k}(p_1, p_2, \dots, p_k, q) + gr_{L_0, L_k, L_{k+1}, \dots, L_{n-1}}(q, p_{k+1}, \dots, p_{n-1}, p_n). \end{aligned}$$

- iii. *$gr_{L_{n-1}, L_{n-2}, \dots, L_0}(p_{n-1}, p_{n-2}, \dots, p_1, p_n) = 2 - n - gr_{L_0, L_1, \dots, L_{n-1}}(p_1, p_2, \dots, p_{n-1}, p_n)$.*

In particular,

$$gr_{L_0, L_1}(p, r) = gr_{L_0, L_1}(p, q) + gr_{L_0, L_1}(q, r), \quad gr_{L_0, L_1}(p, p) = 0,$$

$$gr_{L_1, L_0}(q, p) = gr_{L_0, L_1}(p, q) = -gr_{L_0, L_1}(q, p),$$

and if $\mathcal{M}_{L_0, L_1}(p, q)$ is non-empty, then $gr_{L_0, L_1}(p, q) = 1$.

Proof. The assumption that the L_k are restricted immersed curves implies that the mod 4 reduction of $\text{Mas}_{\ell_{\text{inst}}}(\gamma_0, \dots, \gamma_{n-1})$ is independent of the choice of paths γ_k and therefore $gr_{L_0, L_1, \dots, L_{n-1}}(p_1, \dots, p_n)$ is well defined.

The remaining assertions follow from their counterparts in Proposition 2.9. \square

The function $gr_{L_0, L_1} : (L_0 \cap L_1)^2 \rightarrow \mathbb{Z}/4$ is called *the relative $\mathbb{Z}/4$ grading on $C(L_0, L_1)$* . Proposition 3.8 and Equation (5) imply that the differential (if defined) $\partial : C(L_0, L_1) \rightarrow C(L_0, L_1)$ lowers the relative grading by 1, i.e., $(C(L_0, L_1), \partial)$ is a chain (rather than a cochain) complex.

We thank Matt Hogancamp for formulating the following corollary. Its proof follows quickly from Proposition 3.8, and we omit it.

Corollary 3.9. *Given an ordered list (L_0, \dots, L_{n-1}) of pairwise transverse restricted immersed curves,*

$$gr_{L_0, L_1, \dots, L_{n-1}}(p_1, \dots, p_n) - gr_{L_0, L_1, \dots, L_{n-1}}(q_1, \dots, q_n) = \sum_{k=1}^n gr_{L_{k-1}, L_k}(p_k, q_k).$$

Moreover, if there exists a Maslov index k_p immersed n -gon through (p_1, \dots, p_n) and a Maslov index k_q immersed n -gon through (q_1, \dots, q_n) , then

$$gr_{L_0, L_{n-1}}(p_n, q_n) = k_q - k_p + \sum_{k=1}^{n-1} gr_{L_{k-1}, L_k}(p_k, q_k)$$

□

In the next lemma, we provide a practical formula for gr_{L_0, L_1} in terms of the slope 1 line field and the reversed paths $\alpha_k(t) = \gamma_k(1-t)$. We find this formula to be the simplest to remember, and most of the subsequent calculations of relative gradings in this paper are obtained using this formula, without referring back to Maslov index definitions and conventions. The omitted proof consists applying Equation (2) and Proposition 2.9 to the difference class z .

Lemma 3.10. *The relative $\mathbb{Z}/4$ grading on $C(L_0, L_1)$ is given as follows.*

Let p, q be intersection points of L_0 with L_1 , let α_0 be a path in L_0 from p to q , let α_1 be a path in L_1 from q to p , $\tau(L_0, L_1, \ell_1)_p$ and $\tau(L_0, L_1, \ell_1)_q$ the triple indices with respect to the slope 1 line field ℓ_1 . Then

(15)

$$gr_{L_0, L_1}(p, q) = \mu(L_0, \ell_1)_{\alpha_0} + \mu(L_1, \ell_1)_{\alpha_1} + \tau(L_0, L_1, \ell_1)_p - \tau(L_0, L_1, \ell_1)_q + z(L_0(\alpha_0) * L_1(\alpha_1)).$$

□

When the order is clear from context, we write $gr(p, q)$ rather than $gr_{L_0, L_1}(p, q)$ for the relative $\mathbb{Z}/4$ grading on $C(L_0, L_1)$.

3.8. Finiteness of bigons. When (L_0, L_1) is a transverse pair of restricted immersed curves, we have constructed a relative $\mathbb{Z}/4$ grading on the vector space $C(L_0, L_1)$ spanned by the intersection points of $L_0 : R_0 \rightarrow P^*$ and $L_1 : R_1 \rightarrow P$. To show that $C(L_0, L_1)$ is a chain complex, we must show that $\mathcal{M}(p, q) = \mathcal{M}_{L_0, L_1}(p, q)$ is finite for any pair intersection points p, q . To this end we introduce the notion of an admissible pair.

Definition 3.11. A pair

$$(L_0 : R_0 \rightarrow P, L_1 : R_1 \rightarrow P)$$

of restricted immersed curves in P is called an *admissible pair* provided:

- i. at least one of L_0 or L_1 is a restricted immersed circle,
- ii. if $\alpha_0 : S^1 \rightarrow R_0$ and $\alpha_1 : S^1 \rightarrow R_1$ are loops so that $L_0 \circ \alpha_0$ and $L_1 \circ \alpha_1$ are freely homotopic, then both α_0 and α_1 are nullhomotopic (this holds automatically if one of L_0, L_1 is a restricted immersed arc, since restricted immersed circles are essential),
- iii. L_0 and L_1 intersect transversely.

If we put a complete hyperbolic metric on P^* , then the second assumption is equivalent to the requirement that the unique geodesic representatives of the homotopy classes of L_0 and L_1 are transverse.

Given an admissible pair of restricted curves L_0, L_1 , to each element of $u \in \pi_2(p, q)$ one can assign a *local degree* function f_u , which is an integer valued function with domain the set of complementary regions of $L_0 \cup L_1$, i.e., the path components of $P^* \setminus (L_0 \cup L_1)$. Its value on

a complementary region is the signed number of preimages of a regular value of any smooth representative of u .

Lemma 3.12. *For each pair (p, q) of intersection points, $\pi_2(p, q)$ is either empty or contains a unique element.*

Proof. Since $\pi_2(p, q)$ is either empty or else $\pi_2(p, p)$ acts transitively on $\pi_2(p, q)$, it suffices to prove that $\pi_2(p, p)$ contains a unique class, namely the class of the constant map.

Write $p = (p_0, p_1) \in R_0 \times R_1$. Given $(u, (\gamma_0, \gamma_1)) \in \pi_2(p, p)$, γ_0 (resp. γ_1) is a loop in R_0 (resp. R_1) based at p_0 (resp. p_1). If R_1 is an arc, then $L_1 \circ \gamma_1$ is homotopic rel endpoints to the constant path, and hence so is γ_0 . If R_1 is a circle, then the second assumption of Definition 3.11 implies that γ_0 and γ_1 are nullhomotopic loops. Either way, the images $L_0 \circ \gamma_0$ and $L_1 \circ \gamma_1$ in P^* are homotopic loops based at p .

By the homotopy extension property, $(u, (\gamma_0, \gamma_1))$ may be homotoped in $\pi_2(p, p)$ so that γ_0 and γ_1 are constant. But then u sends the entire boundary of the bigon to p , and hence represents a class in $\pi_2(P^*) = 0$. Thus we may further homotop u rel boundary to the constant map and so $\pi_2(p, p) = 0$, as desired. \square

Corollary 3.13. *Given an admissible pair (L_0, L_1) , each set $\mathcal{M}(p, q)$ is either empty or contains one equivalence class of bigons.*

Proof. Fix intersection points p and q , and suppose that $\mathcal{M}(p, q)$ is non-empty. Choose two immersed bigons u, u' from p to q . Lemma 3.12 implies that $\pi_2(p, q)$ contains a unique element, so their local degree functions are equal. Moreover, since u (and u') are immersed by an orientation preserving immersion, $f_u = f_{u'}$ takes only non-negative values.

Standard arguments now show that u and u' can be reconstructed from the data of their local degrees up to reparameterization, so that u and u' are equivalent. For example, see [29, Theorem 6.8], whose proof applies verbatim to our setting by passing to a compact simply connected submanifold of the universal cover of P^* . \square

Remark 3.14. That $\mathcal{M}(p, q)$ is finite (which is all we require for the assertions in the present article) when L_0 and L_1 are self-transverse immersions can be shown even more easily, as follows. Label the closure of the complementary regions of $P^* \setminus (\text{image}(L_0) \cup \text{image}(L_1))$ by A_1, \dots, A_m . Notice that the boundary of each A_i is a union of arcs $\alpha_{i,j}$ meeting at convex double points. The set of all such arcs, $\{\alpha_{i,j}\}$, can be partitioned into pairs which map to the same arc in P^* , and hence each pair comes with an identification so that the surface obtained by identifying these two arcs immerses into P^* .

If $u \in \pi_2(p, q)$ has all local degrees $f_u(A_i)$ non-negative, take $f_u(A_i)$ copies of A_i , $i = 1, \dots, m$, and label the corresponding edges as $\alpha_{i,j;k}$, $k = 1, \dots, f_u(A_i)$. There are finitely many ways of pairing all the arcs $\{\alpha_{i,j;k}\}$ and gluing them to get a surface which immerses each copy of A_i to its corresponding complementary region. Any immersed bigon from p to q must be equivalent to one of these resulting glued surfaces, hence there are finitely many bigons.

From Theorem 2.14, Proposition 3.8, and Corollary 3.13 we conclude the following.

Theorem 3.15. *If (L_0, L_1) is an admissible pair, then $(C(L_0, L_1), \partial)$ is a relatively $\mathbb{Z}/4$ graded chain complex with \mathbb{F}_2 coefficients.* \square

Definition 3.16. Call the $\mathbb{Z}/4$ graded homology of $(C(L_0, L_1), \partial)$ the *Lagrangian-Floer homology of the (L_0, L_1)* and denote it by $HF(L_0, L_1)$.

3.9. Example: Calculation of $HF(L_0, L_1)$ for $L_0 = L_0^{\epsilon, 0}$ and L_1 the restricted immersed circle of Figure 1. The pair (L_0, L_1) is admissible. In Figure 8, the eight intersection points of the Lagrangian L_1 of Figure 1 with $L_0^{\epsilon, 0}$ are labeled p, q, r, s, t, u, v, w . There is a bigon from p to q and hence $gr(p, q) = 1$. Similarly, there is a bigon from w to v and hence $gr(w, v) = 1$.

We compute $gr(q, s) = 1$ in detail next. Let α_0 be the path in L_0 starting at q , heading down and to the left, around the lower left corner, then back up to s . Let α_1 be the short arc on L_1 from s back to q . Then $\mu(L_0, \ell_1)_{\alpha_0} = 1$ (there is one tangency at the upper right part of the figure, near $(0, 2\pi)$), $\mu(L_1, \ell_1)_{\alpha_1} = 0$ since the arc α_1 is everywhere transverse to the slope 1 line field ℓ_1 . Next, $\tau(L_0, L_1, \ell_1)$ equals 0 at q and 1 at s , and $z(L_0(\alpha_0) * L_1(\alpha_1)) = 1$ since the loop $L_0(\alpha_0) * L_1(\alpha_1)$ goes once around the lower left corner counterclockwise. Using Equation (15) we conclude:

$$gr(q, s) = 1 + 0 + 0 - 1 + 1 = 1.$$

An identical argument gives $gr(t, p) = 1$, $gr(v, r) = 1$, and $gr(u, w) = 1$.

One more calculation is required to complete the calculation of the relative grading, for example $gr(t, r)$. Take α_0 to be the path in L_0 from t to r which heads down and to the left, around the bottom left corner clockwise, then back up to t . Take α_1 the path in L_1 from r back to t which starts by heading to the right, then down and continuing along L_1 until it returns to t . Then $\mu(L_0, \ell_1)_{\alpha_0} = 1$, $\mu(L_1, \ell_1)_{\alpha_1} = 1$, $\tau(L_0, L_1, \ell_1)_t = 0$, $\tau(L_0, L_1, \ell_1)_r = 1$, and $z(L_0(\alpha_0) * L_1(\alpha_1)) = 0$. Thus

$$gr(t, r) = 1 + 1 + 0 - 1 + 0 = 1.$$

From these calculations and additivity of the relative grading, we conclude that $gr(p, r) = 0$, $gr(p, q) = gr(p, u) = 1$, $gr(p, s) = gr(p, w) = 2$, $gr(p, t) = gr(p, v) = 3$. Hence $C(L_0, L_1)$ has rank 2 in each grading. There are only two Maslov index 1 bigons, and hence the differential is given by $\partial p = q$ and $\partial v = w$, and so the homology has rank 1 in each grading.

We introduce a bit of notation that will simplify our descriptions of the $\mathbb{Z}/4$ gradings. The notation (n_0, n_1, n_2, n_3) with n_i non-negative integers denotes the $\mathbb{Z}/4$ graded vector space (over \mathbb{F}_2) whose dimension on grading i is n_i . Thus, for this example,


$$C(L_0, L_1) = (2, 2, 2, 2) \text{ and } HF(L_0, L_1) = (1, 1, 1, 1).$$

4. HOMOTOPY INVARIANCE

We show that the relatively $\mathbb{Z}/4$ graded group $HF(L_0, L_1)$ depends only on the homotopy classes of L_0 and L_1 (rel boundary) in P^* . Our argument follows the approach taken in [1, Proposition 4.1] and presumably will be somewhat familiar to experts. It is worth noting that we make no requirement that the curves be related by a Hamiltonian isotopy.

Theorem 4.1. *Let (L_0, L_1) and (L'_0, L'_1) be two admissible pairs which satisfy:*

- i. *the admissible circles L_0, L'_0 are freely homotopic.*

FIGURE 8. The eight intersection points of L_0 and L_1 generating $C(L_0, L_1)$.

- ii. If L_1, L'_1 are immersed restricted circles they are freely homotopic. If L_1, L'_1 are immersed restricted arcs, they are homotopic rel endpoints. Assume also that near their endpoints, L_1 and L'_1 intersect only at their endpoint.
- iii. L_0, L_1, L'_0, L'_1 are pairwise transverse and have no triple points.

Then

$$HF(L_0, L_1) \cong HF(L'_0, L'_1).$$

as relatively $\mathbb{Z}/4$ graded \mathbb{F}_2 vector spaces.

Proof. In order to avoid the proliferation of sub and superscripts, we make the following notational changes. Set

$$A := L_0, B := L_1, C := L'_0, D := L'_1.$$

And we must show that

$$HF(A, B) = HF(C, D).$$

Consider A and C as immersions of the unit circle $A, C : S^1 \rightarrow P^*$. Also, consider B and D as immersions of the real line \mathbb{R} to P^* , with the understanding that if R_1 is a circle, then B and D are 2π periodic and, if R_1 is an arc, then we identify the interior of R_1 (which maps by B, D to P^*) with \mathbb{R} . In brief, B and D are immersions of \mathbb{R} to P^* which are periodic if R_1 is a circle and proper if R_1 is an arc. The second condition in the hypotheses implies that outside some compact set in \mathbb{R} , B and D are disjoint embeddings, but with the same limit points at $\pm\infty$.

Let $x = A(1)$. The immersion A induces a homomorphism on fundamental groups. Consider the infinite cyclic subgroup

$$Z = \text{Image } A_{\#} : \pi_1(S^1, 1) \rightarrow \pi_1(P^*, x)$$

and let

$$f : (\Sigma, \hat{x}) \rightarrow (P^*, x)$$

denote the (non-regular) cover corresponding to Z . Thus $A : S^1 \rightarrow P^*$ lifts to $\hat{A} : S^1 \rightarrow \Sigma$, with $\hat{A}(1) = \hat{x}$.

Since $\hat{A} : S^1 \rightarrow \Sigma$ generates $\pi_1(\Sigma, \hat{x}) = Z \cong \mathbb{Z}$, the preimage of \hat{A} in the universal cover $\tilde{P}^* \cong \mathbb{R}^2$ is connected, in fact the image of an immersion $\tilde{A} : \mathbb{R} \rightarrow \mathbb{R}^2$. Since $A = L_0$ is unobstructed, \hat{A} is an embedding, from which it follows that $\hat{A} : S^1 \rightarrow \Sigma$ is an embedding. (In the following, we use the notation $\hat{A}, \hat{B}, \hat{C}, \hat{D}$ for lifts of A, B, C to Σ , and $\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}$ for lifts to the universal cover \mathbb{R}^2 .)

It can be easily shown, for example using elementary hyperbolic geometry, that Σ is diffeomorphic to the cylinder $S^1 \times \mathbb{R}$. Fix such a diffeomorphism and the corresponding cover

$$(S^1 \times \mathbb{R}, \hat{x}) \rightarrow (P^*, x).$$

Let $F : S^1 \times [0, 1] \rightarrow P^*$ be a homotopy from A to C . Let $\hat{F} : S^1 \times [0, 1] \rightarrow S^1 \times \mathbb{R}$ be the unique lift of F satisfying $\hat{F}(1, 0) = \hat{x}$. Let $\hat{C}(z) = \hat{F}(z, 1)$. Then \hat{C} is a lift of C to $S^1 \times \mathbb{R}$, and, as with \hat{A} , \hat{C} is an embedding. In particular, since \hat{A} and \hat{C} are homotopic embedded curves in $S^1 \times \mathbb{R}$, \hat{A} and \hat{C} are isotopic.

The following three lemmas will complete the proof of Theorem 4.1.

Lemma 4.2. *If \hat{A} and \hat{C} meet transversely in precisely two points, then $HF(A, B) \cong HF(C, B)$ and $HF(A, D) \cong HF(C, D)$ as relatively $\mathbb{Z}/4$ graded \mathbb{F}_2 vector spaces.*

Lemma 4.3. *There exists a sequence A_0, A_1, \dots, A_r of homotopic restricted immersed circles so that $A_0 = A$, $A_r = C$, and \hat{A}_k intersects \hat{A}_{k+1} transversely in two points.*

Lemma 4.4. *$HF(A, B) \cong HF(A, D)$ as relatively $\mathbb{Z}/4$ graded \mathbb{F}_2 vector spaces.*

Proof of Lemma 4.2. Up to diffeomorphism of the cylinder $S^1 \times \mathbb{R}$, the curves \hat{A} and \hat{C} are illustrated in Figure 9. Also illustrated is a third curve \hat{A}' which meets \hat{A} transversely in two points. We assume that \hat{A}' is very (C^1) close to \hat{A} , so that the preimage of B is also transverse to \hat{A}' and induces a bijection between the intersection points. Three pairs of intersection points $\hat{a}, \hat{b} \in \hat{A} \cap \hat{B}$, $\hat{f}, \hat{c} \in \hat{A}' \cap \hat{C}$, and $\hat{b}, \hat{d} \in \hat{A} \cap \hat{A}'$ are illustrated. Define A' to be the image of \hat{A}' under the covering map $S^1 \times \mathbb{R} \rightarrow P^*$, and let a, b, \dots, f denote the images of $\hat{a}, \hat{b}, \dots, \hat{f}$ in P^* .

We now summarize some facts about the maps μ_2 and μ_3 , defined in Definition 2.13, that will be used to complete Lemma 4.2.

Lemma 4.5. *Consider b as a generator of $C(A, C)$, e as a generator of $C(A, A')$ and f as a generator of $C(C, A')$. Then:*

- i. *For any $x \in C \cap B$ and $y \in A \cap B$, the set $\mathcal{M}_{A, C, B}(b, x, y)$ is finite.*
- ii. *For any $x \in A' \cap B$ and $y \in C \cap B$, the set $\mathcal{M}_{C, A', B}(f, x, y)$ is finite.*
- iii. *For any $x \in A' \cap B$ and $y \in A \cap B$, the set $\mathcal{M}_{A, A', B}(e, x, y)$ is finite.*




FIGURE 9.

iv. For any $x \in A' \cap B$ and $y \in A \cap B$, the set $\mathcal{M}_{A,C,A',B}(b, f, x, y)$ is finite.

Hence the maps $\mu_2(b, -) : C(C, B) \rightarrow C(A, B)$, $\mu_2(f, -) : C(A', B) \rightarrow C(C, B)$, $\mu_2(e, -) : C(A', B) \rightarrow C(A, B)$ and $\mu_3(b, f, -) : C(A', B) \rightarrow C(A, B)$ are well defined.

Proof. For the first statement, first note that to each class $\phi \in \pi_2(b, x, y)$ one can assign an integer local degree to each complementary region of $A \cup C \cup B$, i.e., to each path component of $P^* \setminus (A \cup C \cup B)$ (see the proof of Corollary 3.13).

The homotopy group $\pi_2(b, b)$ corresponding to the pair A, C acts on $\pi_2(b, x, y)$ corresponding to the ordered triple (A, C, B) by attaching a bigon from b to b to the vertex of a triangle with vertices b, x, y at b . We show this action is transitive. From Figure 9 one sees that there exists $\tau_1 \in \pi_2(b, b)$ whose two boundary loops represent generators of $\pi_1(A)$ and $\pi_1(C)$. Let τ_n denote the n th power of τ_1 .

Suppose that $\phi_1, \phi_2 \in \pi_2(b, x, y)$. Denote the image of ϕ_k in $\pi_1(b, x, y)$ by $(\alpha_k, \gamma_k, \beta_k)$, $k = 1, 2$, where α_k is a path in A from y to b , γ_k is a path in C from b to x and β_k is a path in B from x to y .

The loop $\gamma_1 * \gamma_2^{-1}$ in C based at b represents some multiple of the generator of $\pi_1(C)$. Hence, by replacing ϕ_2 by $\tau_n \cdot \phi_2$ for the appropriate n , one may assume that $\gamma_1 * \gamma_2^{-1}$ is nullhomotopic. Using the homotopy extension property we may arrange that $\gamma_1 = \gamma_2$.

The triangles ϕ_1 and ϕ_2 then glue together along γ_1 and γ_2 to provide a free homotopy of the loop $\beta_1 * \beta_2^{-1}$ in B to the loop $\alpha_1 * \alpha_2^{-1}$ in B . Since A, B form an admissible pair (Definition 3.11) it follows that $\alpha_1 * \alpha_2^{-1}$ and $\beta_1 * \beta_2^{-1}$ are nullhomotopic, so that α_1, α_2 (resp. β_1, β_2) are homotopic rel endpoints. We may therefore replace ϕ_2 by another map in the same homotopy class so that $\alpha_1 = \alpha_2$ and $\beta_1 = \beta_2$. Gluing ϕ_1 to ϕ_2 along the boundary arcs α_k, β_k and γ_k yields a class in $\pi_2(P^*) = 0$. Hence $\phi_1 = \phi_2$. We have shown that $\pi_2(b, b)$ acts transitively on $\pi_2(b, x, y)$.

A class $\tau \in \pi_2(b, b)$ determines local degrees for each complementary region of $A \cup C$, and hence also for each complementary region of $A \cup C \cup B$. Lifting to the cylinder, one sees that $\pi_2(b, b) \cong \pi_2(\hat{b}, \hat{b}) \cong \mathbb{Z}$, with $n \in \mathbb{Z}$ corresponding to the class $\tau_n \in \pi_2(\hat{b}, \hat{b})$, whose local

degrees in $S^1 \times \mathbb{R} \setminus (A \cup C)$ are n and $-n$ in the two bounded regions of $S^1 \times \mathbb{R} \setminus (\hat{A} \cup \hat{C})$, and 0 in both unbounded regions. Let W_1 and W_3 denote the two bounded regions in $S^1 \times \mathbb{R} \setminus (\hat{A} \cup \hat{C})$, and W_2 and W_4 denote the two unbounded regions, indexed so that moving clockwise around \hat{b} they are ordered W_1, W_2, W_3, W_4 and so that the local degrees of τ_n are $n, 0, -n, 0$. (see Figure 10, with $n = n_1$ and $n_2 = 0$.)

If $U \subset P^*$ is a small evenly covered disc neighborhood of b , only finitely many of the components of the preimage of U in $S^1 \times \mathbb{R}$ meet W_1 and W_3 . Suppose that K_1 such components meet W_1 and K_2 meet W_3 . Let $K = K_1 - K_2$. Then in P^* , the local degrees of τ_n about b are, in clockwise order, $n + Kn, Kn, -n + Kn, Kn$.

Fix $\phi \in \pi_2(y, x, b)$ and let d_1, d_2, d_3, d_4 denote local degrees of ϕ in the four quadrants around b . The local degrees of $\tau_n \cdot \phi$ are just the sum of the local degrees of τ_n and ϕ . Thus the local degrees of $\tau_n \cdot \phi$ around b are, in clockwise order, $d_1 + n + Kn, d_2 + Kn, d_3 - n + Kn, d_4 + Kn$.

On the other hand, if $\tau_n \cdot \phi$ is represented by a Maslov index 0 immersed 3-gon, then all its corners are convex (Proposition 2.11) and so the local degrees near b must take the form $r + 1, r, r, r$ moving clockwise around b , for some non-negative integer r . In particular,

$$|d_1 - d_3 + 2n| = |(d_1 + n + Kn) - (d_3 - n + Kn)| \leq 1,$$

and so at most two of the classes $\tau_n \cdot \phi$ support Maslov index 0 immersed 3-gons. Each such class determines a finite number of immersed 3-gons, by the same argument given in Remark 3.14. Thus $\mathcal{M}(b, x, y)$ is finite.


The proofs of the second and third assertions are the same, and we leave them to the reader.

The last assertion has a similar proof, so we outline it, highlighting the differences. The ordered 4-tuple (A, C, A', B) gives rise to the sets $\mathcal{M}(b, f, x, y)$ and $\pi_2(b, f, x, y)$. Since A, C, A' are homotopic and A, B form an admissible pair, a similar argument to that used in the triangle case shows that any two classes $\phi_1, \phi_2 \in \pi_2(b, f, x, y)$ are related by the action of $\pi_2(b, b) \times \pi_2(f, f)$: one first lets $\pi_2(b, b)$ act on ϕ_1 to make the boundary paths of ϕ_1 and ϕ_2 along A agree, then let $\pi_2(f, f)$ act to make the boundary paths along C agree. Gluing the two rectangles along these edges yields a twice punctured sphere, giving a free homotopy from a loop in A' to a loop in B , and since (A', B) forms an admissible pair, these loops are both nullhomotopic. The argument then proceeds as in the 3-gon case to conclude that there exists $\tau_{n_1} \in \pi_2(b, b)$ and $\rho_{n_2} \in \pi_2(f, f)$ so that $\tau_{n_2} \cdot \rho_{n_1} \cdot \phi_1 = \phi_2$.

Fix $\phi \in \pi_2(b, f, x, y)$ and assume its local multiplicities in the four quadrants clockwise around b are (d_1, d_2, d_3, d_4) . In the four quadrants near \hat{b} , the multiplicities of $\tau_{n_1} \in \pi_1(b, b)$ are, in clockwise order, $n_1, 0, -n_1, 0$ and the multiplicities of $\rho_{n_2} \in \pi_2(f, f)$ are $n_2, n_2, 0, 0$. Figure 10 illustrates the contributions of τ_{n_1} and ρ_{n_2} to the local multiplicities near b .

As in the triangle case, the local multiplicities of $\tau_{n_2} \cdot \rho_{n_1} \cdot \phi$ near b are, in clockwise order, $d_1 + n_1 + n_2 + K, d_2 + K, d_3 - n_1 + n_2 + K, d_4 + K$, where K is a function of n_1 and n_2 which takes into account how many of the preimages of b in $S^1 \times \mathbb{R}$ lie in the complementary regions on which τ_{n_2} and ρ_{n_1} are supported. But to represent an orientation preserving immersed 4-gon of Maslov index -1 , all corners are convex (Proposition 2.11), and hence these numbers must be a cyclic permutation of $(r + 1, r, r, r)$ for some non-negative integer r . This implies that

$$|d_1 - d_3 + 2n_1| \leq 1 \text{ and } |d_1 + n_1 + n_2 - d_4| \leq 1,$$

FIGURE 10. The local multiplicities of ρ_{n_2} and τ_{n_1} around b

which is only possible for at most four choices of n_1, n_2 . Thus only finitely many classes in $\pi_2(b, f, x, y)$ support immersed Maslov index -1 rectangles, and as in Remark 3.14, this implies that $\mathcal{M}(b, f, x, y)$ is finite. \square

Continuing the proof of Lemma 4.2. Consider b as a generator of $C(A, C)$. Any bigon from b to another intersection point of A with C has a unique lift to the cover $S^1 \times \mathbb{R}$, and hence must be a bigon from b to a . From Figure 9 one sees that there are precisely 2 such bigons up to equivalence, and hence $\partial b = 2a = 0$ (we are using \mathbb{F}_2 coefficients). Thus b is a cycle. Similar arguments show that $e \in C(A, A')$ and $f \in C(C, A')$ are also cycles.

The map μ_2 satisfies the A_2 relation, which, applied to b and an arbitrary $x \in C(C, B)$ (writing ∂ instead of μ_1) says:

$$0 = \mu_2(\partial b, x) + \mu_2(b, \partial x) + \partial \mu_2(b, x).$$

Since $\partial b = 0$, this implies that $\mu_2(b, -) : C(C, B) \rightarrow C(A, B)$ is a chain map.

The cycle $f \in C(C, A')$ determines a chain map $\mu_2(f, -) : C(A', B) \rightarrow C(C, B)$. The product $\mu_2(b, f) \in C(A, A')$ equals e , since immersed triangles lift to the cover, and there is precisely one oriented immersed Maslov index zero triangle for the ordered triple (A, C, A') from (\hat{b}, \hat{f}) to \hat{x} for an intersection point \hat{x} of \hat{A} and \hat{A}' , namely the embedded triangle with $\hat{x} = \hat{e}$. Similarly, the cycle $e = \mu_2(b, f)$ determines a chain map

$$\mu_2(e, -) : C(A', B) \rightarrow C(A, B).$$

The A_3 relation gives, for an arbitrary generator x of $C(A', B)$:

$$\begin{aligned} 0 &= \mu_3(\partial b, f, x) + \mu_3(b, \partial f, x) + \mu_3(b, f, \partial x) + \mu_2(\mu_2(b, f), x) + \mu_2(b, \mu_2(f, x)) + \partial \mu_3(b, f, x) \\ &= H(\partial x) + \mu_2(e, x) + \mu_2(b, -) \circ \mu_2(f, x) + \partial H(x) \end{aligned}$$

where $H : C(A', B) \rightarrow C(A, B)$ is defined by $H(x) = \mu_3(b, f, x)$. In other words, H is a chain homotopy from the composite

$$C(A', B) \xrightarrow{\mu_2(f, -)} C(C, B) \xrightarrow{\mu_2(b, -)} C(A, B)$$

to

$$\mu_2(e, -) : C(A', B) \rightarrow C(A, B).$$

Now $\mu_2(e, -)$ is a chain isomorphism: this is an immediate consequence of the fact that \hat{A} and \hat{A}' are C^1 close, so that each intersection point p' of A' with B corresponds to precisely one intersection point p of A with B , the correspondence induced by a unique Maslov index 0 immersed triangle associated to the ordered triple (A, A', B) From (e, p') to p . This proves that the map $\mu_2(b, -) : C(C, B) \rightarrow C(A, B)$ induces a surjection $HF(C, B) \rightarrow HF(A, B)$, for any B .

The fact that $\mu_2(b, -)$ induces an injection $HF(C, B) \rightarrow HF(A, B)$ for any B is proved by a very similar argument as surjection, using the immersed curves illustrated in Figure 11. In this case, \hat{C}' is a curve C^1 close to C , and one shows that $z \in C(C', A)$ is a cycle, $\mu_2(z, b) = w \in C(C', C)$, and the composite of chain maps

$$C(C, B) \xrightarrow{\mu_2(b, -)} C(A, B) \xrightarrow{\mu_2(z, -)} C(C', B)$$

is chain homotopic to the chain isomorphism $\mu_2(w, -) : C(C, B) \rightarrow C(C', B)$. This proves $\mu_2(b, -)$ induces an injection $HF(C, B) \rightarrow HF(A, B)$ for any B .




FIGURE 11.

Finally we show that the relative $\mathbb{Z}/4$ grading is preserved. Suppose that $x_1, x_2 \in C(C, B)$ and $\mu_2(b, x_1) = \sum_i m_i y_i$ and $\mu_2(b, x_2) = \sum_j n_j z_j$ in $C(A, B)$, with m_i, n_j non-zero. Hence there exist Maslov index zero immersed 3-gons for the ordered triple (A, C, B) from (b, x_1) to y_i and from (b, x_2) to z_j .

Then Corollary 3.9 gives

$$gr_{A,B}(y_i, z_j) = gr_{C,B}(x_1, x_2) + gr_{A,C}(b, b) = gr_{B,C}(x_1, x_2).$$

This says that the chain map $\mu_2(b, -) : C(C, B) \rightarrow C(A, B)$ preserves the relative $\mathbb{Z}/4$ grading, and hence also the induced map on homology. The same argument shows the chain maps $\mu_2(f, -)$ and $\mu_2(e, -)$ preserve the relative grading. Thus, $\mu_2(b, -)$ induces an isomorphism $HF(A, B) \rightarrow HF(A, C)$ of relatively $\mathbb{Z}/4$ graded vector spaces. \square

Proof of Lemma 4.3. If \hat{A} and \hat{C} intersect non-trivially (hence in an even number of points), then the existence of the sequence $\hat{A}_0 = \hat{A}, \hat{A}_1, \dots, \hat{A}_r = \hat{C}$ with the stated property is an immediate consequence of Lemma 4.2 of [1]. If \hat{A} and \hat{C} are disjoint, then one can take a parallel copy of \hat{A} and perform a Reidemeister 2 move that introduces a pair of intersection points with \hat{A} and a pair with \hat{B} . \square

Proof of Lemma 4.4. If B and D are homotopic restricted immersed circles, then the proof follows exactly along the same lines as the proof of Lemmas 4.2 and 4.3, reversing the roles (and order) of A, C and B, D .

If B and D are homotopic restricted immersed arcs, then a different proof is needed. It is convenient to put a complete hyperbolic metric on P^* and to let $\mathbb{H} \rightarrow P^*$ denote the universal cover. Let $\tilde{B} : \mathbb{R} \rightarrow \mathbb{H}$ be a lift of B . Since D is homotopic rel endpoints to B , there is a lift $\tilde{D} : \mathbb{R} \rightarrow \mathbb{H}$ of D with the same limit points at the circle at infinity. The assumption that the limiting slopes at the endpoints are distinct and that B and D are transverse imply that the closures of \tilde{B} and \tilde{D} in the closed disk $\overline{\mathbb{H}}$ intersect in finitely many points.

If \tilde{B} and \tilde{D} are disjoint, then their closures bound a bigon in $\overline{\mathbb{H}}$ with vertices on the circle at infinity. If \tilde{B} and \tilde{D} intersect in one point, then clearly there are a pair of bigons with boundary in their closures, each including one point on the boundary of $\overline{\mathbb{H}}$. Finally, if $\tilde{B} \cap \tilde{D}$ consists of more than one point, then the result of D.B.A Epstein [13, Lemma 3.2] (see also [6, Lemma A.10]) shows that one can find an embedded bigon in \mathbb{H} for \tilde{B} and \tilde{D} .

In each of these three cases, one can find a sequence of embedded arcs $\tilde{B} = \tilde{B}_0, \tilde{B}_1, \dots, \tilde{B}_r = \tilde{D}$ in \mathbb{H} so that \tilde{B}_k intersects \tilde{B}_{k+1} transversely in one interior point and in their endpoints. The argument is illustrated in Figure 12. The figure on the left corresponds to the first case, when $\tilde{B} \cap \tilde{D} = \emptyset$. The figure on the right corresponds to the third case, where an interior bigon between \tilde{B} and \tilde{D} is used to construct an arc \tilde{B}_1 which intersects $\tilde{B}_0 = \tilde{B}$ in one interior point and \tilde{D} in two fewer points, providing the required induction step. If \tilde{B} and \tilde{D} started out with an odd number of intersection points in their interior, one eventually reaches \tilde{B}_r which meets \tilde{D} in one point in their interior, and if they started out with an even number of intersection points in their interior, one reaches \tilde{B}_r whose interior misses \tilde{D} , in which case one can add one more step as in the first case.




FIGURE 12.

FIGURE 13.

The counterpart to Lemma 4.5 is much simpler in the case of arcs. Consider the ordered triple (A, D, B) . If $p \in A \cap D$, $q \in D \cap B$, and $r \in A \cap B$, and then $\pi_2(p, q, r)$ is either empty or contains a single class. In fact, if $(u_i, (\alpha_i, \gamma_i, \beta_i)) \in \pi_2(p, q, r)$, $i = 1, 2$, then since B and D are arcs we may use the homotopy extension property to assume $\beta_1 = \beta_2$ and $\gamma_1 = \gamma_2$. Then gluing u_1 to u_2 along α_i and β_i shows that $\alpha_1 \alpha_2^{-1}$ is nullhomotopic, so that we may assume further than $\gamma_1 = \gamma_2$. Since $\pi_2(P^*) = 0$, it follows that $u_1 = u_2$ in $\pi_2(p, q, r)$. As before, this implies that $\mathcal{M}(p, q, r)$ is finite, and hence $\mu_2 : C(A, D) \times C(D, B) \rightarrow C(A, B)$ is defined. The same argument applies to the triple (A, D, D') to show that $\mu_2 : C(A, D) \times C(D, D') \rightarrow C(A, D')$ is defined, and to the 4-tuple (A, D, B, D') to show that $\mu_3 : C(A, D) \times C(D, B) \times C(B, D') \rightarrow C(A, D')$ is defined.

Using the same notation for points in the universal cover and their images in P^* , we see that $\partial x = 0$ in $C(D, B)$, hence the A_2 relation shows that $\mu_2(-, x) : C(A, D) \rightarrow C(A, B)$ is a chain map.

Observe that $\partial(y+z) = 2w = 0$ in $C(B, D')$. Also, $e+f = \mu_2(x, y+z)$ in $C(D, D')$. Furthermore, $\partial(e+f) = 2u = 0$. Hence $\mu_2(-, e+f) : C(A, D) \rightarrow C(A, D')$ is a chain map. In fact, $\mu_2(-, e+f)$ is a chain isomorphism, since \tilde{D} is close to \tilde{D}' and A misses neighborhoods of $\pm\infty$, so there is a unique intersection point w' of A with D' for each intersection point w of A with D . If w lies between e and u , then $\mu_2(w, e) = w'$ and $\mu_2(w, f) = 0$, and if w lies between x and f , then $\mu_2(w, f) = w'$ and $\mu_2(w, e) = 0$.

Substituting these calculations into the A_3 relation

$$\begin{aligned} 0 &= \mu_3(\partial w, x, y+z) + \mu_3(w, \partial x, y+z) + \mu_3(w, x, \partial(y+z)) \\ &\quad + \mu_2(\mu_2(w, x), y+z) + \mu_2(w, \mu_2(x, y+z)) + \partial\mu_3(w, x, y+z) \end{aligned}$$

and defining $H : C(A, D) \rightarrow C(A, D')$ by $H(w) = \mu_3(w, x, y+z)$ yields

$$0 = H\partial(w) + \mu_2(\mu_2(w, x), y+z) + \mu_2(w, e+f) + \partial H(w).$$

So that the chain isomorphism $\mu_2(-, e+f) : C(A, D) \rightarrow C(A, D')$ is chain homotopic to the composite $C(A, D) \xrightarrow{\mu_2(-, x)} C(A, B) \xrightarrow{\mu_2(-, y+z)} C(A, D')$. Hence the chain map $\mu_2(-, x) : C(A, D) \rightarrow C(A, B)$ is injective for all restricted immersed circles A so that (A, B) and (A, D) are admissible.

The reader can safely be left the task of showing that $\mu_2(-, x) : C(A, D) \rightarrow C(A, B)$ is injective for all restricted immersed circles A , by producing an embedded \tilde{B}' close to \tilde{B} and constructing a right inverse to $\mu_2(-, x)$. This is done by analogy with the symmetry between Figures 9 and 11. The fact that the relative $\mathbb{Z}/4$ gradings is preserved is proved as before.

This completes the proof of Lemma 4.4, and hence also the proof of Theorem 4.1. \square

Corollary 4.6. *$HF(L_0^{\epsilon, g}, L_1)$ is independent of $\epsilon > 0$, the function g and the homotopy class of L_1 .* \square

5. CALCULUS

In this section, we make four technical but useful observations which streamline the calculation of $HF(L_0, L_1)$ when $L_0 = L_0^{\epsilon, g}$. These calculations demonstrate the ease of working with the slope 1 line field ℓ_1 and Equation (15).

5.1. We show $gr(x_+, x_-) = 1$ when x is a transverse intersection of L_1 with the diagonal arc, and x_+, x_- the two corresponding intersection points with $L_0 = L_0^{\epsilon, g}$ for $\epsilon > 0$ and g small, as indicated in Figure 14. We take α_0 to be the path in L_0 starting at x_+ which heads down and to the left, around the bottom left corner, and back up to x_- . The path α_1 from x_- back to x_+ is the short path which contains the diagonal intersection point x .

In this case, $\mu(\alpha_0, \ell_1) = 1$, $\mu(\alpha_1, \ell_1) = 0$, $\tau(L_0, L_1, \ell_1)_{x_+} = 0$, $\tau(L_0, L_1, \ell_1)_{x_-} = 1$, and $z(\alpha_0 * \alpha_1) = 1$ yielding $gr(x_+, x_-) = 1$.

5.2. Suppose that L_1 intersects the diagonal arc Δ transversely and that $L_0 = L_0^{\epsilon, g}$ for $\epsilon > 0$ and g small. Suppose p, q are intersection points between L_0 and Δ . Let p_+, q_+ be the corresponding intersection points with the part of L_0 in the front of P which has slope slightly less than 1, as indicated in Figure 14.

Then the arc α_0 along L_0 from p_+ to q_+ can be chosen to lie entirely on the front face of the pillowcase, and has slope slightly less than 1, hence $\mu(\alpha_0, \ell_1) = 0$. Moreover, since the slope of ℓ_1 equals 1, L_1 is transverse to Δ , and ϵ and g are small, the triple index terms τ at p_+




FIGURE 14.

and q_+ are both zero. This leaves only the terms $\mu(\alpha_1, \ell_1)$ and $z(\alpha_0 \alpha_1)$ in the formula for the grading. Therefore, *in this situation*

$$(16) \quad gr(p_+, q_+) = \mu(\alpha_1, \ell_1) + z(\alpha_0 * \alpha_1).$$

A similar calculation applies to the other pair p_-, q_- .

5.3. If R is an arc, $L_0 \cap L_1$ may (and in our applications will) contain a distinguished point which may be used to promote the relative grading to an absolute grading. To understand the meaning of the following lemma, the reader should locate the distinguished point r_+^ϵ in the examples illustrated below in Figures 15, 21, 22, and 23.

Lemma 5.1. *Suppose that $L_1 : R \rightarrow P$ is a restricted immersed arc and one of the endpoints (which we denote by r_+), maps to the corner $(0, 0)$ of P with limiting slope bounded away from 1. Then for all small ϵ , there is a unique continuously varying intersection point r_+^ϵ of L_0^ϵ and L_1 satisfying $\lim_{\epsilon \rightarrow 0} r_+^\epsilon = r_+$.*

Proof. This follows simply from the requirement that limiting slope bounded away from 1. \square

If L_1 satisfies the hypothesis of Lemma 5.1, then, given the additional data of a choice of $\sigma \in \mathbb{Z}/4$, one can endow $HF(L_0, L_1)$ with the absolute $\mathbb{Z}/4$ grading which places r_+^ϵ in grading σ .

In our applications below, L_1 will be associated to a knot K in a homology 3-sphere. In this setting the hypotheses of Lemma 5.1 hold and we take σ to be the signature of the knot K .

5.4. A special class of restricted immersed circles, which we call vertically monotonic, arise in many of our examples and have particularly simple Lagrangian-Floer homology.

Definition 5.2. Suppose that $L_1 : R \rightarrow P^*$ is a restricted immersed circle, with domain parameterized by $[0, 2\pi]$, and let $\tilde{L}_1 = (\gamma(t), \theta(t)) : [0, 2\pi] \rightarrow \mathbb{R}^2$ denote its lift to the branched cover (8).

- Call $L_1 : R \rightarrow P^*$ *vertically monotonic* if \tilde{L}_1 misses the vertical line segments $\gamma = k\pi, k \in \mathbb{Z}$, and if its tangent slope satisfies $|\frac{d}{dt} \tilde{L}_1| > 1$. Thus L_1 winds around the

pillowcase without intersecting the left or right edges, and is everywhere transverse to the line field ℓ_1 (as well as the slope -1 line field).

- Define the *vertical degree* of L_1 to be the absolute value of the difference of the vertical coordinates of $\tilde{L}_1(2\pi)$ and $\tilde{L}_1(0)$. Thus

$$d = \frac{1}{2\pi} |\theta(0) - \theta(2\pi)|.$$

Notice that the vertical degree is a homotopy invariant. In particular, it is well defined for any circle homotopic to a vertically monotonic circle.

Proposition 5.3. *Suppose that $L_1 : R \rightarrow P^*$ is a vertically monotonic restricted immersed circle. Then the vertical degree d of L_1 is even. Moreover, for ϵ small enough, all differentials are zero and $HF(L_0, L_1)$ has rank $\frac{d}{2}$ in each of the 4 grading degrees.*

Proof. Since $L_1 : R \rightarrow P^*$ is vertically monotonic, it is transverse to the line field ℓ_1 and hence $\mu(L_1(R), \ell_1) = 0$. Moreover, $L_0^{\epsilon, g}$ is transverse to L_1 for all small enough ϵ, g . Fix a transverse $L_0 = L_0^{\epsilon, g}$ with ϵ, g small.

Since $\tilde{L}_1 : [0, 2\pi] \rightarrow \mathbb{R}^2$ misses the vertical line segments $\gamma = k\pi$, $z(L_1(R))$ is equal to twice the vertical degree d of L_1 . Since $z(L_1(R)) \equiv -\mu(L_1(R)) \pmod{4}$, $2d \equiv 0 \pmod{4}$ so that d is even.

Let p, q be intersection points of L_0 and L_1 and suppose that there were a bigon (u, α_0, α_1) from p to q . The bigon misses the corners of P and hence lifts to a bigon $(\tilde{u}, \tilde{\alpha}_0, \tilde{\alpha}_1)$ in \mathbb{R}^2 with one edge along the preimage of L_1 and one along the preimage of L_0 .

For $\epsilon > 0$ sufficiently small, the connected components of the preimage in \mathbb{R}^2 of L_0 are very close to lines of slope 1 through $(0, k\pi)$, and hence $\tilde{\alpha}_0$ is nearly a straight segment of slope 1. On the other hand, since the tangent lines to \tilde{L}_1 have slope bounded away from 1, the lift $\tilde{\alpha}_1$, which starts at $\tilde{\alpha}_0(1)$, cannot terminate at $\tilde{\alpha}_0(0)$, contradicting the fact that $\tilde{\alpha}_0$ and $\tilde{\alpha}_1$ bound a bigon. Hence, for sufficiently small ϵ , there are no bigons, so that all differentials are zero. The circle $L_1(R)$ winds monotonically around P , intersecting the diagonal arc Δ in d points, and hence intersecting L_0 in $2d$ points, so that $HF(L_0, L_1)$ has rank $2d$.

It remains to calculate the relative gradings. Since L_1 is transverse to ℓ_1 , Equation 16 shows that this reduces to calculating z . One vertical wind around P encircles two corner points and therefore changes z by 2 mod 4. Hence the generators come in $\frac{d}{2}$ pairs, and, as explained in Section 5.1 and indicated in Figure 14, alternate between contributing in (relative) gradings 0, 1 and 2, 3. This completes the argument. \square

Proposition 5.3 can be strengthened by combining it with Corollary 4.6 as follows.

Theorem 5.4. *Suppose that $L_1 : R \rightarrow P^*$ is a restricted immersed circle which misses the left and right edges of the pillowcase. Then the vertical degree d is defined and even, and for any $\epsilon > 0$ and g , $HF(L_0^{\epsilon, g}, L_1)$ has rank $\frac{d}{2}$ in each of the 4 grading degrees.*

Proof. Since L_1 misses the right and left edges of the pillowcase, it is homotopic to a vertically monotonic immersed circle. The proof is therefore a consequence of Corollary 4.6. \square

6. TRACELESS REPRESENTATION VARIETIES OF 2-STRANDED TANGLES

Hereafter, a *restricted immersed 1-manifold* is a disjoint union of a single restricted immersed arc and a finite number of restricted immersed circles. This prompts us to make the

following changes in our notation. The first immersed curve, $L_0 = L_0^{\epsilon, g} : S^1 \rightarrow P$ henceforth has domain the circle, and the domain of L_1 has multiple components, so we now use the symbol R_0 to denote the arc component for L_1 . More precisely, suppose R is the disjoint union of an arc R_0 and finitely many circles R_1, \dots, R_n . $L_1 : R \rightarrow P$ a map so that the restriction of L_1 to R_0 is a restricted immersed arc and the restriction to each R_i , $i \geq 1$ is a restricted immersed circle. The Lagrangian-Floer homology $H^\natural(L_0, L_1)$ is defined to be the direct sum of the homologies $HF(L_0^{\epsilon, g}, L_1|_{R_i})$, where ϵ, g are chosen so that $(L_0^{\epsilon, g}, L_1|_{R_i})$ form an admissible pair for each i .

Each summand is relatively $\mathbb{Z}/4$ graded, although initially there is neither a relative nor absolute $\mathbb{Z}/4$ grading on all of $H^\natural(L_0, L_1)$ if L_1 is not connected. The notation H^\natural is adopted in order to indicate the relationship to the reduced instanton homology IH^\natural .

We now introduce traceless character varieties. We identify $SU(2)$ with the set of unit quaternions throughout. For the basic properties of the unit quaternions and their Lie algebra, we refer the reader to Section 2 of [18]. A *traceless quaternion* means a quaternion with zero real part.

Given a pair (A, B) consisting of a (compact) manifold and a properly embedded codimension 2 submanifold, call a representation $\pi_1(A \setminus B) \rightarrow SU(2)$ *traceless* if it sends all meridians of B to the conjugacy class $C(\mathbf{i})$ of \mathbf{i} . Define the *traceless character variety* (or traceless flat moduli space)

$$R(A, B) = \{\rho : \pi_1(A \setminus B) \rightarrow SU(2) \mid \rho \text{ is traceless}\} / \text{conjugation}$$

An embedding of pairs $(A_1, B_1) \subset (A_2, B_2)$ induces a restriction map $R(A_2, B_2) \rightarrow R(A_1, B_1)$.

Consider a decomposition of a pair (X, K) , where K is a knot (or link) in a homology 3-sphere X , and X contains a separating 2-sphere $S \subset X$ which intersects K transversally in four points. We assume that one of the two regions S bounds is a 3-ball D , and that $D \cap K$ is a standard trivial 2-stranded tangle.

$$(17) \quad (X, K) = (Y, T) \cup_{(S, \{a, b, c, d\})} (D, U)$$

We refer to (17) as a *2-tangle decomposition associated to the knot* (X, K) . We fix an identification $D = B^3$ so that

$$U = \left\{ \left(\pm \frac{1}{\sqrt{2}}, 0, t \right) \mid t \in \left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right] \right\}$$

and fix an identification $\partial(D, U) = (S^2, \{a, b, c, d\})$ with $a = \left(\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}} \right)$, $b = \left(-\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}} \right)$, $c = \left(-\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}} \right)$, $d = \left(\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}} \right)$. We call (Y, T) a *2-tangle associated to the knot* (X, K) .

Observe that to recover (X, K) from (Y, T) requires only a choice of identification

$$\iota : (Y, T) \cong (S^2, \{a, b, c, d\})$$

(in the same way that a Dehn filling is determined by a manifold with torus boundary and an identification of its boundary with the boundary of a solid torus). We will omit the choice of ι from the notation since the identification will be clear from context.

To a 2-tangle decomposition of (X, K) , one associates the diagram

$$(18) \quad \begin{array}{ccc} & R(S, \{a, b, c, d\}) & \\ L_1 \nearrow & & \nwarrow L_0 \\ R_\pi(Y, T) & & R_{\pi_\epsilon}^{\natural}(D, U) \\ & \nwarrow & \nearrow \\ & R_{\pi''}^{\natural}(X, K) & \end{array}$$

where π, π_ϵ refer to certain *holonomy perturbations* $\pi'' = \pi \cup \pi_\epsilon$, and $R_\pi(M, L)$ denotes the corresponding π -*perturbed traceless character variety*.

Holonomy perturbations are used to make the Chern-Simons function (whose Morse theory defines instanton homology) have only non-degenerate critical points. They are constructed and explained in the context of the traceless character varieties in [18, Section 7] and also in Section 8.2 below. They were introduced in gauge theory by Donaldson, Floer, Taubes, and others [10, 32, 16].

It is not necessary for this article to understand precisely what the \natural superscript means beyond knowing the statement of Theorem 6.2 below. But roughly, $R_{\pi_\epsilon}^{\natural}(D, U)$ refers to $R_{\pi_\epsilon}(D, U \cup E)$, where E is an additional small meridian component to one of the components of U and one considers representations which come from flat connections on an $SO(3)$ bundle with w_2 dual to an arc spanning U and E . This construction, introduced by Kronheimer-Mrowka in [21], is an ingredient in the definition of reduced instanton knot homology. We refer to [21, 18] for the details.

The space $R(S^2, \{a, b, c, d\})$ is a pillowcase. Indeed the following simple proposition is proved in [18] (and elsewhere). In the statement we abuse notation and let a, b, c, d also denote the oriented meridians of the four punctures.

Proposition 6.1 ([18], Proposition 3.1). *There is a surjective quotient map*

$$\psi : \mathbb{R}^2 \rightarrow R(S^2, \{a, b, c, d\})$$

given by

$$\psi(\gamma, \theta) : a \mapsto \mathbf{i}, \quad b \mapsto e^{\gamma \mathbf{k}} \mathbf{i}, \quad c \mapsto e^{\theta \mathbf{k}} \mathbf{i}, \quad d \mapsto e^{(\theta - \gamma) \mathbf{k}} \mathbf{i}.$$

The map ψ induces a homeomorphism of the pillowcase P with $R(S^2, \{a, b, c, d\})$. The four corner points are the image under ψ of the lattice $(\pi\mathbb{Z})^2$, and correspond to abelian non-central representations. All other points correspond to non-abelian representations. \square

We urge the reader not to confuse $(S^2, \{a, b, c, d\})$ with the pillowcase $P = R(S^2, \{a, b, c, d\})$, a homeomorphic space!

The only fact we will need to recall about $R_{\pi_\epsilon}^{\natural}(D, U)$ is the following, which follows immediately by combining [18, Theorem 7.1] with Theorem 9.1, proved below.

Theorem 6.2. *Given any $\epsilon > 0$ and $g \in \mathcal{X}$, there is a holonomy perturbation π_ϵ depending on ϵ and g so that $R_{\pi_\epsilon}^{\natural}(D, U)$ is a circle, and the restriction to the pillowcase (the northwest map in Diagram (18)) is given by a map $L_0^{\epsilon, g}$ of Definition 3.4.* \square

What Proposition 6.1 and Theorem 6.2 tell us is that a decomposition of a knot or link into two 2-tangles, one of which is trivial, gives the pillowcase P and a map $L_0 : S^1 \rightarrow P^*$.

The remaining input needed to define a Lagrangian-Floer homology as in Section 2 is provided by the 1-manifold $R = R(Y, T)$ and $L_1 : R \rightarrow P$ the restriction map $R(Y, T) \rightarrow R(S^2, \{a, b, c, d\})$, as indicated in Diagram (18).

Loosely speaking, $L_1 : R(Y, T) \rightarrow R(S^2, \{a, b, c, d\})$ is generically a union of a restricted immersed arc and some number of restricted immersed circles. The arc arises as one component of the space of traceless binary dihedral representations of $\pi_1(Y \setminus T)$, and the endpoints correspond to the two conjugacy classes of abelian traceless representations (Theorem 3.2 of [17].)

It is not always literally true, however, that $L_1 : R(Y, T) \rightarrow P$ is a restricted immersed 1-manifold. It is true for certain tangles associated to 2-bridge knots [18, Section 10], and for some, but not all torus knots [17].

In fact, there exist decompositions of knots for which $R(Y, T)$, rather than being a smooth 1-manifold, is instead a singular real algebraic variety of dimension greater than or equal to 1. For example in [18, Section 11] (see Figure 19) it is shown that for a tangle associated to the $(3, 4)$ torus knot, $R(Y, T)$ is a singular 1-dimensional variety, homeomorphic to the letter ϕ . Many more examples are given in [17]. In general one can construct examples so that $R(Y, T)$ is highly singular and has strata of high dimension by placing local knots in one of the strands of a 2-tangle. Hence the traceless character varieties $R(Y, T)$ must first be desingularized before we can apply the construction of the Lagrangian-Floer theory in the pillowcase. In order to preserve the relationship to gauge theory and instanton homology, we use holonomy perturbations to smooth $R(Y, T)$.

The space $R(Y, T)$ for a certain natural 2-tangle decomposition of a torus knot is typically singular ([18, 17]). We prove below that any torus knot admits a 2-tangle decomposition and an arbitrarily small holonomy perturbation π so that $R_\pi(Y, T)$ is a compact 1-manifold with two boundary points, and L_1 an immersion which satisfies all the requirements to be a restricted immersed 1-manifold except possibly the requirement that it have no fishtails. Based on index calculations and examples, it is reasonable to expect that for any knot, arbitrarily small holonomy perturbations exist which make $L_1 : R_\pi(Y, T) \rightarrow P$ a restricted immersed 1-manifold.

Conjecture 6.3. *For any 2-tangle (Y, T) in the 3-ball (or a homology 3-ball), there exist arbitrarily small holonomy perturbations π so that $R_\pi(Y, T)$ is a compact 1-manifold with two boundary points and the restriction map $L_1 : R_\pi(Y, T) \rightarrow R(S^2, \{a, b, c, d\})$ is a restricted immersed 1-manifold on each component in the sense of Definition 3.6.*

Given a 2-tangle decomposition of a knot and a perturbation π which satisfies the conclusion of Conjecture 6.3, denote by $H^\natural(Y, T, \pi)$ the resulting Lagrangian-Floer homology of the complex $C(L_0, L_1)$. We will simplify this to $H^\natural(Y, T)$ if the perturbation π is clear from context.

As explained in [18], if $\epsilon \neq 0$ is small and $L_0 = L_0^{\epsilon, 0}$ intersects L_1 transversely, then the intersection points of L_0 and L_1 also form generators of the reduced instanton homology $I^\natural(X, K)$. Theorem 9.1 implies that this holds for $L_0 = L_0^{\epsilon, g}$ for any small $g \in \mathcal{X}$.

We state this formally.

Proposition 6.4. *Given a small perturbation π which makes $L_1 : R_\pi(Y, T) \rightarrow P$ a restricted immersed 1-manifold and a transverse $L_0 = L_0^{\epsilon, g}$ with ϵ and g small, there is a (possibly different) differential*

$$\partial_{KM} : C(L_0, L_1) \rightarrow C(L_0, L_1)$$

so that the homology of $(C(L_0, L_1), \partial_{KM})$ is the reduced instanton homology $I^\natural(X, K)$. \square

The differential ∂_{KM} is defined by Kronheimer-Mrowka in terms of singular instantons on cylinders $(X, K) \times \mathbb{R}$. There is a well known procedure for producing approximate instantons from bigons in character varieties associated to lagrangian intersection diagrams; see for example [34, Section 4]. It is therefore not unreasonable to conjecture that there is a relationship between $I^\natural(X, K)$ and $H^\natural(Y, T, \pi)$. Indeed, we have found these to be isomorphic in every example we have computed. We extend Conjecture 6.3 to an optimistic “Atiyah-Floer” type conjecture:

Conjecture 6.5. *Given a knot (X, K) in a homology 3-sphere, there exists a 2-tangle decomposition as in Equation (17), such that for suitably small generic perturbations π , $L_1 : R_\pi(Y, T) \rightarrow P$ is a restricted immersed 1-manifold and $H^\natural(Y, T, \pi)$ is isomorphic to the reduced instanton homology $I^\natural(X, K)$.*

In the remainder of this article we establish some partial results and carry out calculations which provide evidence for these conjectures. The reader should realize, however, that there are no non-zero differentials in $C(L_0, L_1)$ between generators which lie on different path components of L_1 . We know of no reason why this should be true for the instanton complex. It is likely that there are differentials in the instanton complex which don’t appear in $C(L_0, L_1)$. For example, the pairs of generators p_+, p_- near each intersection point p of L_1 with the diagonal arc Δ , described in Section 5.1, arise from a holonomy perturbation which “tilts” a Bott-Morse circle of critical points of the Chern-Simons function [18]. Analogy with finite-dimensional Morse theory suggests that there exists a cancelling pair of gradient flow lines (i.e., instantons) from p_+ to p_- in the Kronheimer-Mrowka instanton complex, whereas there are no bigons connecting these points of intersection.

6.1. Absolute grading. We remark that, by construction, $H^\natural(Y, T, \pi)$ splits as the direct sum over the path components R_0, R_1, \dots, R_n of $R(Y, T)$:

$$H^\natural(Y, T, \pi) = \oplus_i H^\natural(L_0, L_1(R_i)).$$

and that each of the summands admits a relative $\mathbb{Z}/4$ grading. The relative grading of the summand corresponding to the arc component R_0 can be promoted to an absolute $\mathbb{Z}/4$ grading for small perturbations, using the knot signature, as follows.

Assume that $(X, K) := (Y, T) \cup_\iota (D, U)$ is a 2-tangle decomposition of a knot K in an integer homology sphere X . The signature of K , $\sigma(K)$, is an even integer. There are two traceless abelian representations of $\pi_1(Y \setminus T)$, r_+ and r_- distinguished by the property that r_+ extends to $\pi_1(X \setminus K)$ (and r_- does not). The point r_+ is a Morse critical point of the Chern-Simons function, and a regular point of $R(X, K)$. In particular, it remains regular after small perturbations.

The points r_+ and r_- are endpoints of an embedded arc of binary dihedral representations, which, by Theorem 3.2 of [17], is the image in the pillowcase under the branched cover (8) of an embedded linear segment joining two lattice points. This line segment has slope different

from 1 (the slope is different from 1 since the 2-fold branched cover of a knot in a homology sphere is a rational homology sphere, so the integer $h(bc^{-1})$ in Theorem 3.2 of [17] is non-zero). In particular, the arc of binary dihedral representations is properly immersed (in fact, embedded) in P .

Small perturbations only change the slopes near the endpoints slightly, and one can keep them bounded away from 1. By Lemma 5.1 there is a unique intersection point r_+^ϵ of $R(Y, T)$ and $L_0^{\epsilon, g}$ for all small ϵ, g . We promote the relative grading of the subcomplex corresponding to the component R_0 by declaring

$$(19) \quad gr(r_+^\epsilon) = \sigma(K)$$

for small perturbations.

We have not found an elementary approach to promote the relative grading of the generators of the subcomplexes associated to the circle components $R_i, i > 0$, and so we will use the following awkward definition as a consequence of Proposition 6.4: choose a generator on each circle component and declare its absolute grading to be the one assigned to it by Kronheimer-Mrowka in [23, Proposition 4.4].

A proof that the relative $\mathbb{Z}/4$ grading of generators of $C(L_0, L_1)$ (Definition 3.7) coincides with the grading assigned to these generators (by Proposition 6.4) of singular instanton knot homology by [23, Proposition 4.4] is given by using splitting theorems for spectral flow [25, 9, 5]. We outline how this is done, referring to [5] for details.

First, the relative grading is defined to be the mod 4 reduction of the spectral flow of the Hessian of the Chern-Simons function (acting on singular connections) along a path joining a pair a_0, a_1 of critical points, i.e., perturbed flat connections. If the restrictions of a_0, a_1 to $Y \setminus T$ can be joined by a smooth path of flat connections, i.e., by a smooth path in $R_\pi(Y, T)$, then the approach of [5, Theorem 3.9] can be modified to show that the spectral flow equals the Maslov index along the path of the tangent space of the immersed 1-manifold $R_\pi(Y, T) \rightarrow P^*$ in the pillowcase, with respect to some *a priori* unknown line field ℓ_{inst} , and hence is given as in Definition 3.7.

Changing the homotopy class of a line field determines a difference class $z \in H^1(P^*; \mathbb{Z}/4)$. The identification of ℓ_{inst} is therefore equivalent to the identification of z . Its identification with the explicit class of Definition 3.1 is completed by calculating a few examples of 2-bridge knots, whose instanton homology is known, to deduce the values of z on a basis of 1-cycles in $H_1(P^*)$.

This argument, combined with the additivity of spectral flow under composition of paths of self-adjoint operators, also shows that if $L_1 : R_i \rightarrow P^*$ is an immersion of a smooth circle component $R_i \subset R_\pi(Y, T)$, then L_1 satisfies the condition $\mu(L_1(R_i), \ell_{\text{inst}}) \equiv 0 \pmod{4}$ required of restricted immersed circles. In this case, one uses the fact that the two smooth paths in R_i joining a_0 to a_1 must give the same relative $\mathbb{Z}/4$ grading, since the relative grading in the singular instanton complex is well defined (and independent of the tangle decomposition). As the proof of Proposition 3.8 shows, this is only possible if $L_1 : R_i \rightarrow P^*$ satisfies $\mu(L_1(R_i), \ell_{\text{inst}}) \equiv 0 \pmod{4}$.

7. EXAMPLES: 2-BRIDGE KNOTS

Two-bridge knots can be described as the union of two trivial tangles along a 4-punctured sphere. We recall some of the results about their tangle decompositions from [18]. In particular, we will show that for such a tangle decompositions of a 2-bridge knot K , $L_1 : R(Y, T) \rightarrow P$ is a restricted immersed (in fact linearly embedded) arc which meets $L_0^{\epsilon, 0}$ transversely in $\det(K)$ points, and that all differentials in the Lagrangian-Floer complex are zero.

These facts, together with the identification of the relative $\mathbb{Z}/4$ gradings in $C(L_0, L_1)$ and the singular instanton complex via a spectral flow splitting theorem as explained above, imply that $H^\natural(Y, T)$ is isomorphic to the reduced instanton homology $I^\natural(S^3, K)$, which is known [23] to equal the reduced Khovanov homology $Kh^{red}(K^m)$ of the mirror of K for a 2-bridge knot. We conjectured in [18] that placing the distinguished generator r_+^ϵ in grading $\sigma(K)$ agrees with Kronheimer-Mrowka's absolute grading [23, Proposition 4.4] (a conjecture borne out in all our calculations) and, modulo this point, for 2-bridge knots, $H^\natural(Y, T)$, $I^\natural(S^3, K)$, and $Kh^{red}(K^m)$ (with its bigrading (i, j) reduced to $i - j + 1 \bmod 4$) contain the same information. In particular, Conjecture 6.5 holds for 2-bridge knots.

Suppose that (p, q) are relatively prime integers, with p odd and positive. Then there is a 2-bridge knot $K = K(p, q)$ determined by the condition that its 2-fold branched cover is the lens space $L(p, q)$. In [18, Figure 13], a 2-tangle decomposition $(S^3, K) = (Y, T) \cup (D, U)$ determined by a continued fraction expansion of $\frac{p}{q}$ is described. It is proved that $R(Y, T)$ is a smooth arc and the restriction map to the pillowcase is given, in \mathbb{R}^2 coordinates, by

$$R(Y, T) \cong [0, \pi] \ni t \mapsto (qt, (q - p)t) \in P.$$

Thus $L_1 : R(Y, T) \rightarrow P$ is a restricted embedded arc. In particular, no perturbation π is needed to smooth $R(Y, T)$. Hence we can choose $L_0 = L_0^{\epsilon, 0}$ for a small ϵ and form the chain complex $C(L_0, L_1)$. Since L_1 maps in linearly and L_0 is close to the linear arc Δ , there can be no immersed bigons, and therefore all differentials are zero.

There are p intersection points of L_1 with L_0 . In fact, there are $\frac{p+1}{2}$ intersection points of $(\gamma(t), \theta(t)) = (qt, (q - p)t)$, $t \in [0, \pi]$ with the arc Δ , these occur at

$$(20) \quad x_\ell = \left(q \frac{2\pi\ell}{p}, (q - p) \frac{2\pi\ell}{p} \right), \quad \ell = 0, 1, \dots, \frac{p-1}{2}.$$

The points x_ℓ , $\ell > 0$, each give rise to a pair of intersection points x_ℓ^+, x_ℓ^- of L_1 with L_0^ϵ , and the point x_0 gives rise to the distinguished point r_+^ϵ of Lemma 5.1. Hence $C(L_0, L_1)$ and $H^\natural(Y, T)$ have rank p . The intersection points are illustrated in the case of $K = K(11, -5)$ (7_2 in the knot tables) in Figure 15.

We show how to calculate the gradings. First, the observation of Section 5.1 shows that $gr(x_\ell^+, x_\ell^-) = 1$. Next, recall that we promote the relative grading to an absolute grading by setting $gr(r_+^\epsilon) = \sigma(K) \bmod 4$, where $\sigma(K)$ denotes the signature of the knot K . Thus the determination of all other gradings is reduced to calculating $gr(r_+^\epsilon, x_\ell^+)$ for $\ell = 1, \dots, p$.

The slope $\frac{q-p}{q}$ is not equal to ± 1 since p and q are relatively prime. There are four different cases to be considered, depending on the slope. For simplicity we assume $\frac{q-p}{q}$ is positive and greater than 1; the other cases are treated similarly.

For each $x_\ell^+, \ell = 1, \dots, p$, one can find a path α_0 in L_0 from r_+^ϵ to x_ℓ^+ which lies on the front of the pillowcase. One can then take a path α_1 in L_1 from x_ℓ^+ back to r_+^ϵ . Notice that since L_1 is an arc, the path α_1 is unique.

We are in the situation explained in Section 5.2 and can calculate gradings using the Equation (16). Since the tangent line to L_1 is everywhere transverse to the slope one line field ℓ_1 , Equation (16) simplifies further to $gr(r_+^\epsilon, x_\ell^+) = z(\alpha_0\alpha_1)$.

The same formula holds when $\frac{q-p}{q} < -1$. When $-1 < \frac{q-p}{q} < 1$, an entirely similar calculation yields $gr(r_+^\epsilon, x_\ell^-) = z(\alpha_0\alpha_1)$.

We summarize:

Theorem 7.1. *Let $K = K(p, q) \subset S^3$ be a 2-bridge knot with $p > 0$ odd, and equip it with the 2-tangle decomposition described in [18, Figure 13]. Then $L_1 : R(Y, T) \rightarrow P$ is a linearly embedded arc of slope $\frac{q-p}{q}$, and hence a restricted immersed arc. Taking $L_0 = L_0^{\epsilon, 0}$ with $\epsilon > 0$ small, $C(L_0, L_1)$ has rank p , generated by the points r_+^ϵ and $x_\ell^\pm, \ell = 1, \dots, \frac{p-1}{2}$. The $\mathbb{Z}/4$ grading is determined by*

$$gr(r_+^\epsilon) = \sigma(K), \quad gr(x_\ell^+, x_\ell^-) = 1$$

and, letting x_ℓ° denote x_ℓ^+ if $|\frac{q-p}{q}| > 1$ and x_ℓ^- if $|\frac{q-p}{q}| < 1$

$$gr(r_+^\epsilon, x_\ell^\circ) = z(\alpha)$$

where α is the loop in P^* which starts at r_+^ϵ , follows L_0 to x_ℓ° , then returns to r_+^ϵ along L_1 , and $z \in H^1(P^*; \mathbb{Z}/4)$ is the class of Definition 3.1. All differentials are zero and hence $C(L_0, L_1) \cong H^\natural(Y, T)$. \square

The knot $K(11, -5)$ has signature $\sigma = 2$, and hence $gr(r_+^\epsilon) = 2$. The generators are illustrated in Figure 15. The loop α which follows L_0 from r_+^ϵ to x_1^+ on the front of the pillowcase and then follows L_1 back to r_+^ϵ satisfies $z(\alpha) = 2$. Hence $gr(r_+^\epsilon, x_1^+) = 2$. A similar calculation applies to the other x_ℓ^+ and yields

$$gr(r_+^\epsilon, x_2^+) = 1, gr(r_+^\epsilon, x_3^+) = 0, gr(r_+^\epsilon, x_4^+) = 3, \text{ and } gr(r_+^\epsilon, x_5^+) = 2.$$

The gradings $gr(r_+^\epsilon, x_\ell^-)$ are computed using the fact that $gr(x_\ell^+, x_\ell^-) = 1$. Thus, in the notation introduced above, $H^\natural(Y, T) = (3, 2, 3, 3)$.

The choice $(p, q) = (11, 6)$ gives a different tangle decomposition for the same knot $K = K(11, -5) = K(11, 6)$. The resulting homology is again $(3, 2, 3, 3)$.

The choice $(p, q) = (5, -3)$ yields a tangle decomposition of the Figure 8 knot. The map L_1 is illustrated (with different notation) in [18, Figure 16]. There are 5 generators, $r_+^\epsilon, x_1^\pm, x_2^\pm$, and computing gradings using z yields $gr(r_+^\epsilon) = \sigma(K) = 0, gr(x_1^+) = 3, gr(x_2^+) = 2$, and hence $H^\natural = (1, 1, 2, 1)$. This agrees with the calculation of reduced instanton homology and reduced Khovanov homology of the Figure 8 knot. Choosing $(p, q) = (5, 2)$ gives a different tangle decomposition for the Figure 8 knot, but again yields $H^\natural = (1, 1, 2, 1)$.

The trefoil knot corresponds to $(p, q) = (3, -1)$; one calculates $H^\natural = (1, 0, 1, 1)$. The same answer is obtained when taking instead $(p, q) = (3, 2)$.

Theorem 7.1 can easily be used (and implemented in a computer algebra program) to compute $H^\natural(Y, T) \cong I^\natural(S^3, K) \cong Kh^{red}(K^m)$ for any 2-bridge knot K . In particular, this




FIGURE 15. The intersection of L_0 and L_1 in P for the 2-bridge knot $K(11, -5)$.

gives a novel approach to computing the reduced Khovanov homology of 2-bridge knots (with its bigrading (i, j) reduced to $i - j + 1 \bmod 4$).

We point out that the main *new* ingredients contained in this discussion of 2-bridge knots which were not implicit in [18] are first, the construction of the complex $C(L_0, L_1)$ associated to a tangle decomposition of a 2-bridge knot, and second, the use of the cohomological invariant $z \in H^1(P^*; \mathbb{Z}/4)$ and the Maslov index to define and compute the relative grading.

8. SOME GENERAL PROPERTIES OF $R(Y, T)$

8.1. Structure of $R(Y, T)$ near the abelian points. Suppose that (Y, T) is a 2-tangle. Our goal (not fully realized in this article) is to establish Conjecture 6.3. To this end, we start by showing that the two boundary points of $R_\pi(Y, T)$ are well defined for small perturbations, and correspond to the precisely two *abelian* representations in $R(Y, T)$, namely the conjugacy classes of the two representations

$$r_\pm : \pi_1(Y \setminus T) \rightarrow \{\pm 1, \pm \mathbf{i}\} \subset SU(2)$$

uniquely characterized (since $H_1(Y \setminus T) = \mathbb{Z} \oplus \mathbb{Z}$, generated by μ_1, μ_2) by

$$r_\pm(\mu_1) = \mathbf{i}, r_\pm(\mu_2) = \pm \mathbf{i}.$$

The following proposition proves that r_+ and r_- each have a neighborhood in $R(Y, T)$ homeomorphic to a half-open interval. The outline of the argument is as follows: the space $R(Y, T)$ is identified with a subspace of the space of conjugacy classes of representations of the 2-fold branched cover (the *equivariant representations* in the sense of [30]). Then a Kuranishi model argument shows that the representation space is locally a half open interval near the lifts of r_{\pm} .

Proposition 8.1. *For a 2-tangle T in an integer homology ball Y , each of the two abelian traceless representations r_{\pm} has a neighborhood in $R(Y, T)$ homeomorphic to a half-open interval. The restriction map to the pillowcase P properly embeds each half-open interval, taking the endpoints to distinct corners with limiting slope not equal to 1.*

The proof is an extension of [17, Theorem 3.2]. That theorem identifies the subvariety $R^{tbd}(Y, T) \subset R(Y, T)$ of traceless binary dihedral representations with the disjoint union of one arc and a number of circles (the number determined by the torsion submodule of the homology of the 2-fold branched cover of Y branched along T). The endpoints of the arc are precisely r_+ and r_- , and the arc of binary dihedrals is linearly embedded into the pillowcase with slope different from 1 (see Section 6.1). Hence what must be shown is that there are no non-binary dihedral representations in small enough neighborhoods of r_{\pm} .

We begin with a lemma which permits us to transfer the problem to one about the 2-fold branched cover of (Y, T) . To this end, Let $c : \pi_1(Y \setminus T) \rightarrow \{\pm 1\}$ be the unique homomorphism sending both μ_1 and μ_2 to -1 (this is just the homomorphism $r_+^2 = r_-^2$). Let $B \rightarrow Y$ denote the corresponding 2-fold branched cover. Denote the preimage of T by \tilde{T} . Consider $\pi_1(B \setminus \tilde{T})$ as the index 2 subgroup of $\pi_1(Y \setminus T)$, i.e., as the kernel of c . Let $\tilde{\mu}_1, \tilde{\mu}_2$ denote the meridians of the two components of \tilde{T} . Hence, in $\pi_1(Y \setminus T)$, $\tilde{\mu}_i = \mu_i^2$.

Denote by $R_{\pm 1}(B, \tilde{T})$ the space of conjugacy classes of representations of $\pi_1(B \setminus \tilde{T})$ which take the $\tilde{\mu}_i$ to ± 1 . Since the square of a traceless element of $SU(2)$ is -1 , restriction to the index 2 subgroup defines a map

$$R(Y, T) \rightarrow R_{-1}(B, \tilde{T}).$$

Lemma 8.2. *Pointwise multiplication by c defines a $\mathbb{Z}/2$ action on $R(Y, T)$ with fixed points the traceless binary dihedral representations. The restriction map $R(Y, T) \rightarrow R_{-1}(B, \tilde{T})$ is constant on $\mathbb{Z}/2$ orbits and embeds the quotient $R(Y, T)/\mathbb{Z}/2 \subset R_{-1}(B, \tilde{T})$.*

Assuming Lemma 8.2, the proof of Proposition 8.1 can be completed as follows.

Denote by \tilde{r}_{\pm} the restrictions of r_{\pm} to the index 2 subgroup $\pi_1(B \setminus \tilde{T})$. Then \tilde{r}_{\pm} takes values in the center $\{\pm 1\}$ of $SU(2)$ and $\tilde{r}_{\pm}(\tilde{\mu}_i) = r_{\pm}(\mu_i^2) = -1$.

It follows that pointwise multiplication of a representation by \tilde{r}_+ defines a continuous map $R_{-1}(B, \tilde{T}) \rightarrow R_1(B, \tilde{T})$. This map is a homeomorphism (in fact real analytic isomorphism) with inverse given again by multiplication by \tilde{r}_+ .

Let $\chi(B)$ denote the space of conjugacy classes of (all) $SU(2)$ representations of $\pi_1(B)$. The Seifert-Van Kampen theorem shows that the restriction $\chi(B) \rightarrow R_1(B, \tilde{T})$ is a homeomorphism. Hence we have a sequence of maps :

$$R(Y, T) \rightarrow R(Y, T)/\mathbb{Z}/2 \subset R_{-1}(B, \tilde{T}) \cong R_1(B, \tilde{T}) \cong \chi(B).$$

It therefore suffices to prove that a neighborhood of $\tilde{r}_+\tilde{r}_\pm$ in $\chi(B)$ is homeomorphic to a half-open interval. Notice that $\tilde{r}_+\tilde{r}_+ : \pi_1(B) \rightarrow SU(2)$ is the trivial representation, and $\tilde{r}_+\tilde{r}_- : \pi_1(B) \rightarrow SU(2)$ is central but non-trivial (it takes $\mu_1\mu_2$ to -1).

The 3-manifold B has torus boundary and has first homology isomorphic to $\mathbb{Z} \oplus O$ for O an odd torsion abelian group, since T is a tangle in a homology ball (see [17, Section 3] for details).

The Kuranishi method identifies a neighborhood of c_\pm in $R(B)$ with $K^{-1}(0)/SU(2)$, where $K : H^1(B; su(2)_{ad\ c_\pm}) \rightarrow H^2(B; su(2)_{ad\ c_\pm})$. The adjoint action of c_\pm is trivial since c_\pm is central, and hence these are untwisted cohomology groups with coefficients in $su(2) = \mathbb{R}^3$. The universal coefficient theorem gives $H^1(B; \mathbb{R}^3) = \mathbb{R}^3$ and $H^2(B; \mathbb{R}^3) = 0$, so that c_\pm has a neighborhood homeomorphic to $\mathbb{R}^3/SU(2) = \mathbb{R}^3/SO(3) \cong [0, 1)$, as desired.

Proof of Lemma 8.2. First, if ρ represents a conjugacy class in $R(Y, T)$, then the function $c\rho(\gamma) = c(\gamma)\rho(\gamma)$ is again a representation, since c takes values in the center $\{\pm 1\}$. Moreover, since $\ker c = \pi_1(B \setminus \tilde{T})$, the restrictions of ρ and $c\rho$ to $\pi_1(B \setminus \tilde{T})$ agree. Since $c^2 = 1$, this shows that multiplication by c defines a $\mathbb{Z}/2$ action on $R(Y, T)$ and the restriction $R(Y, T) \rightarrow R_1(B, \tilde{T})$ factors through the quotient of this $\mathbb{Z}/2$ action.

Conversely, suppose $\rho_1, \rho_2 : \pi_1(Y \setminus T) \rightarrow SU(2)$ are two traceless representations whose restriction to the index 2 subgroup $\pi_1(B \setminus \tilde{T})$ are equal. For clarity, denote this restriction by $\bar{\rho}$, so $\bar{\rho} = \rho_1|_{\ker c} = \rho_2|_{\ker c}$.

We claim that, perhaps after conjugating ρ_2 without changing its restriction to $\ker c$, $\rho_1(\mu_1)$ and $\rho_2(\mu_1)$ commute. To see this, first note that for each $\tau \in \ker c$,

$$(21) \quad \rho_1(\mu_1\tau\mu_1^{-1}) = \bar{\rho}(\mu_1\tau\mu_1^{-1}) = \rho_2(\mu_1\tau\mu_1^{-1})$$

so that

$$(22) \quad [\rho_2(\mu_1)^{-1}\rho_1(\mu_1), \bar{\rho}(\tau)] = 1 \text{ for all } \tau \in \ker c$$

If $\bar{\rho}$ has non-abelian image, Equation (22) implies that $\rho_2(\mu_1)^{-1}\rho_1(\mu_1)$ is central, so that $\rho_2(\mu_1) = \pm\rho_1(\mu_1)$ and hence they commute. If $\bar{\rho}$ has central image, then conjugating ρ_2 by any element of $SU(2)$ does not change its restriction to $\ker c$, and since $\rho_1(\mu_1)$ and $\rho_2(\mu_1)$ are traceless, they are conjugate. Hence ρ_2 can be conjugated so that $\rho_1(\mu) = \rho_2(\mu)$ and their restrictions to $\ker c$ agree.

Consider as a final case that $\bar{\rho}$ has abelian non-central image. We show that again ρ_2 can be conjugated without changing its restriction to $\ker c$ to make $\rho_1(\mu_1)$ and $\rho_2(\mu_1)$ commute. Choose a traceless quaternion \mathbf{q} so that the image of $\bar{\rho}$ lies in the circle subgroup $\mathbf{S} := \{e^{\theta\mathbf{q}}\}$. Then Equation (22) shows that $\rho_2(\mu_1)^{-1}\rho_1(\mu_1)$ lies in \mathbf{S} . If one of $\rho_1(\mu_1)$ or $\rho_2(\mu_1)$ lies in \mathbf{S} then they both do since their product does, and hence they commute. Suppose that neither lies in \mathbf{S} . Equation (21) shows that conjugation by $\rho_1(\mu_1)$ and $\rho_2(\mu_1)$ leaves the circle \mathbf{S} invariant. This in turn shows that there exists an element of \mathbf{S} which conjugates $\rho_2(\mu_1)$ to $\rho_1(\mu_1)$. This conjugation leaves \mathbf{S} fixed, so that we have shown that in this final case, ρ_2 can be conjugated without changing its restriction to $\ker c$ to make $\rho_1(\mu_1)$ and $\rho_2(\mu_1)$ commute.

Define $f : \pi_1(Y \setminus T) \rightarrow \{\pm 1\}$ by the formula

$$f(\gamma) = \begin{cases} 1 & \text{if } \gamma \in \ker c \\ \rho_1(\mu_1)\rho_2(\mu_1)^{-1} & \text{if } \gamma \notin \ker c. \end{cases}$$

Then it is easy to see that f is a homomorphism (using the fact that $\rho_1(\mu_1)$ and $\rho_2(\mu_1)$ commute). Moreover, a simple calculation shows that

$$\rho_2(\gamma) = f(\gamma)\rho_1(\gamma) \text{ for all } \gamma \in \pi_1(Y \setminus T).$$

Note that there are exactly two possibilities for f since $\ker c$ has order 2. In fact, the two possibilities are the trivial homomorphism and c . This proves that the restriction map $R(Y, T) \rightarrow R(B, \tilde{T})$ factors through an injective map on the orbit space of this $\mathbb{Z}/2$ action.

It remains to prove that the fixed points are exactly the traceless binary dihedral representations ([17, Definition 3.1]). Suppose that $c\rho$ is conjugate to ρ for $\rho \in R(Y, T)$. Thus there exists $g \in SU(2)$ so that $g\rho(\gamma)g^{-1} = c(\gamma)\rho(\gamma)$ for all $\gamma \in \pi_1(Y \setminus T)$. In particular, $g\rho(\gamma)g^{-1} = \rho(\gamma)$ for all $\gamma \in \ker c$. Since $c(\mu_1) = -1$, $g \neq \pm 1$, so that g lies in a unique circle subgroup which we denote \mathbf{S} .

If ρ sends every $\gamma \in \ker c$ to the center $\{\pm 1\}$, then the image of ρ lies in the subgroup $\{\pm 1, \pm \rho(\mu_1)\}$ of order 4, and ρ is traceless binary dihedral.

On the other hand, if there exists $\gamma \in \ker c$ such that $\rho(\gamma) \neq \pm 1$, then g and $\rho(\gamma)$ commute, and hence ρ sends all of $\ker c$ into \mathbf{S} . Furthermore, for each γ in the non-trivial coset, $\rho(\gamma)^{-1}g\rho(\gamma) = -g$, which implies that $\rho(\gamma)$ is traceless and $\mathbf{S} \cup \rho(\gamma)\mathbf{S}$ is (a conjugate of) the binary dihedral subgroup containing the image of ρ . Hence ρ is traceless binary dihedral. \square

8.2. Perturbations. Proposition 8.1 shows that $R(Y, T)$ is a 1-manifold with boundary near the two abelian representations r_{\pm} . The space $R(Y, T)$ is a real algebraic variety, but in general it may be singular. To prove Conjecture 6.3 for some (Y, T) one must first desingularize $R(Y, T)$.

There are various ways to smooth the singular space $R(Y, T)$; we restrict attention to holonomy perturbations since these have a gauge theoretical counterpart which permits us to compare our constructions to those of [21, 23]. In particular, with this choice of perturbations, Proposition 6.4 identifies the generators of the reduced knot instanton homology chain complex with the intersection points of $R_{\pi}(Y, T)$ and L_0 in the pillowcase for any appropriate perturbation π .

We recall how to understand holonomy perturbations on the level of representations. What follows can be taken as a definition. The reader should keep in mind, however, that the perturbed equations we give below arise from a perturbation of the Chern-Simons functional on the space of traceless $SU(2)$ connections. In particular, what we call a perturbation function is essentially the derivative of the conjugacy invariant function on $SU(2)$ which is used to perturb the Chern-Simons function.

A holonomy perturbation is associated to a pair $\pi = (E, f)$, where

- i. E is an embedding $E : S^1 \times D^2 \subset Y \setminus T$ (we use E also as notation for the image $E(S^1 \times D^2)$), and
- ii. f is a perturbation function, i.e., $f \in \mathcal{X} = \{f \in C^{\infty}(\mathbb{R}, \mathbb{R}) \mid f \text{ is odd, } 2\pi \text{ periodic}\}$.

Call a representation $\rho : \pi_1(Y \setminus (T \cup E)) \rightarrow SU(2)$ a π -perturbed traceless representation if ρ takes the meridians of T to $C(\mathbf{i})$, the conjugacy class of \mathbf{i} , and satisfies the *perturbation condition* on the meridian $\mu_E = E(\{1\} \times \partial D^2)$ and longitude $\lambda_E = E(S^1 \times \{1\})$:

$$(23) \quad \rho(\lambda_E) = e^{\alpha Q} \text{ implies } \rho(\mu_E) = e^{f(\alpha)Q}$$

for $Q \in su(2)$. Then define the *perturbed traceless flat moduli space* $R_\pi(Y, T)$ to be the space of conjugacy classes of π -perturbed traceless representations. We refer the reader to [11, 16, 32]; expositions tailored to our notation can be found in [14, Lemma 61] and [18, Section 7].

More generally, one can choose a collection $E_i, i = 1, \dots, n$ of disjoint embeddings, and corresponding functions f_i define $\pi = \{E_i, f_i\}$, and take $R_\pi(Y, T)$ to be the space of conjugacy classes of π -perturbed traceless representations, defined by requiring the perturbation condition (23) to hold for each i . One useful choice is $f_i(x) = \epsilon_i \sin(x)$ for some small ϵ_i .

The following proposition shows that the two abelian representations are stable with respect to (sup norm of f_i) small perturbations.

Proposition 8.3. *For small enough perturbations, $R_\pi(Y, T)$ contains exactly two conjugacy classes of abelian representations. These are sent to distinct corner points in the pillowcase by the restriction map $R_\pi(Y, T) \rightarrow R(S^2, \{a, b, c, d\})$.*

Proof. Let μ_1, μ_2 denote meridians of the two components of T . Let $\mu_{E_1}, \dots, \mu_{E_n}$ denote the meridians of the perturbation curves. Then $\mu_1, \mu_2, \mu_{E_1}, \dots, \mu_{E_n}$ generate $H_1(Y \setminus T)$.

Let $\ell_i(\mu_1, \mu_2, \mu_{E_1}, \dots, \mu_{E_n}), i = 1, \dots, n$, express the longitude λ_{E_i} in $H_1(Y \setminus T)$ as a linear combination of the meridians of the meridians.

Identify the diagonal maximal torus in $SU(2)$ with the circle S^1 and let $T^{n+2} = (S^1)^{n+2}$. For each $\delta \geq 0$, Let $\pi(\delta)$ denote the perturbation data obtained by multiplying each f_i by δ . Then define a self-map of T^{n+2}

$$Q(\delta) : T^{n+2} \rightarrow T^{n+2}$$

as follows. The first two coordinates encode the traceless condition and are given by

$$Q(\delta)_1(e^{\theta_1 \mathbf{i}}, e^{\theta_2 \mathbf{i}}, e^{\alpha_1 \mathbf{i}}, \dots, e^{\alpha_n \mathbf{i}}) = -e^{2\theta_1 \mathbf{i}}, \quad Q(\delta)_2(e^{\theta_1 \mathbf{i}}, e^{\theta_2 \mathbf{i}}, e^{\alpha_1 \mathbf{i}}, \dots, e^{\alpha_n \mathbf{i}}) = -e^{2\theta_2 \mathbf{i}}.$$

The remaining coordinates encode the perturbation condition:

$$Q(\delta)_{i+2}(e^{\theta_1 \mathbf{i}}, e^{\theta_2 \mathbf{i}}, e^{\alpha_1 \mathbf{i}}, \dots, e^{\alpha_n \mathbf{i}}) = e^{\alpha_i \mathbf{i}} e^{-\delta f_i(\ell_i(\theta_1, \theta_2, \alpha_1, \dots, \alpha_n)) \mathbf{i}}.$$

Then $Q(\delta)^{-1}(1, \dots, 1)$ parameterizes the perturbed traceless abelian representations (not conjugacy classes) with values in the diagonal maximal torus of $SU(2)$, with respect to the functions δf_i : the point $(e^{\theta_1 \mathbf{i}}, e^{\theta_2 \mathbf{i}}, e^{\alpha_1 \mathbf{i}}, \dots, e^{\alpha_n \mathbf{i}}) \in Q(\delta)^{-1}(1, \dots, 1)$ corresponds to the representation

$$\pi_1(B \setminus (T \cup E)) \rightarrow H_1(B \setminus (T \cup E)) \rightarrow S^1 \subset SU(2)$$

sending each meridian to its corresponding coordinate.

The proof is completed by observing that $Q(0)$ is a covering map, hence a submersion. Since submersions are stable, $Q(\delta)^{-1}(1, \dots, 1)$ varies by an isotopy for small δ . \square

For any perturbation π , restricting to the boundary punctured sphere induces a map to the pillowcase

$$R_\pi(Y, T) \rightarrow R(S^2, \{a, b, c, d\}).$$

The two abelian representations guaranteed to persist after small perturbations by Proposition 8.3 necessarily are mapped to corners of the pillowcase, since the restriction of an abelian representation is abelian, and non-corner points are non-abelian, as one can see from Proposition 6.1.

Putting the Propositions 6.1, 8.1 and 8.3, together, we conclude the following.

Theorem 8.4. *Let (Y, T) be a 2-tangle in a \mathbb{Z} -homology ball. Then, for any sufficiently small holonomy perturbation π , there are two abelian perturbed flat representations $r_{\pm} \in R_{\pi}(Y, T)$ with neighborhoods U_{\pm} in $R_{\pi}(Y, T)$ half-open intervals. The restriction map*

$$R_{\pi}(Y, T) \rightarrow R(S^2, \{a, b, c, d\})$$

restricts to an immersion on $U_+ \cup U_-$ which takes r_{\pm} to distinct corners of the pillowcase, with slope $\neq 1$. \square

Thus Theorem 8.4 reduces the problem of defining $H^{\natural}(Y, T)$ for a 2-tangle T to finding an (arbitrarily) small holonomy perturbation π so that

- i. $R_{\pi}(Y, T) \setminus \{r_+, r_-\}$ is a smooth 1-manifold.
- ii. The restriction of $L_1 : R_{\pi}(Y, T) \rightarrow P$ to the arc component is an immersion into P^* containing no fishtails.
- iii. The restriction of $L_1 : R_{\pi}(Y, T) \rightarrow P^*$ to each circle component is an immersion into P^* containing no fishtails.

It is well known that calculations of Zariski tangent spaces using Poincaré-Lefschetz duality show that if $R_{\pi}(Y, T)$ is a smooth 1-manifold away from the two endpoints, then the restriction map $L_1 : R_{\pi}(Y, T) \rightarrow P$ immerses $R_{\pi}(Y, T) \setminus \{r_+, r_-\}$ into P^* . Thus for a given (Y, T) , what is needed is a holonomy perturbation which desingularizes $R(Y, T)$ so that the resulting restriction to P has no fishtails.

9. PERTURBING NEAR THE 2-SPHERE

In this section we construct holonomy perturbations in $(S^2, \{a, b, c, d\}) \times I$ which induce a family of Hamiltonian isotopies of the pillowcase. These were used in Sections 3 and 6 to make L_0 and L_1 transverse. These will also be used for other purposes below and in further work.

Consider the product pair

$$(S^2 \times I, \{a, b, c, d\} \times I).$$

Its traceless character variety is P , and the traceless character variety of its boundary

$$(S^2 \times \{0, 1\}, \{a, b, c, d\} \times \{0, 1\})$$

is $P \times P$. The restriction map

$$R(S^2 \times I, \{a, b, c, d\} \times I) \rightarrow R(S^2 \times \{0, 1\}, \{a, b, c, d\} \times \{0, 1\})$$

is the diagonal map $P \rightarrow P \times P$, which we consider as the graph of the identity map $P \rightarrow P$.

Given suitable perturbation data π for $(S^2 \times I, \{a, b, c, d\} \times I)$, the restriction map

$$R_{\pi}(S^2 \times I, \{a, b, c, d\} \times I) \rightarrow R(S^2 \times \{0, 1\}, \{a, b, c, d\} \times \{0, 1\})$$

gives a *Lagrangian correspondence* $c_{\pi} : P \rightarrow P$. Choosing a path from the trivial perturbation to π gives a homotopy of the identity to c_{π} . We focus on a special class of π for which c_{π} is an explicitly defined diffeomorphism.

Figure 16 shows the 4-punctured 2-sphere with the four based meridian generators a, b, c, d based at a point s . An additional curve e is also indicated.




FIGURE 16.

Let $E : S^1 \times D^2 \rightarrow S^2 \setminus \{a, b, c, d\} \times I$ be a tubular neighborhood of the curve obtained by pushing e into the interior of $S^2 \times I$. Fix a perturbation function $f \in \mathcal{X}$ and let $\delta = (E, f)$ denote the perturbation data. Recall that f can be any smooth odd, 2π periodic function.

Theorem 9.1. *With perturbation data $\delta = (E, f)$, the map*

$$P \rightarrow R_\delta(S^2 \times I, \{a, b, c, d\} \times I)$$

induced by the inclusion $S^2 \times \{0\} \rightarrow S^2 \times I$ is a homeomorphism, and the composite

$$P \rightarrow R_\delta(S^2 \times I, \{a, b, c, d\} \times I) \rightarrow R(S^2 \times \{0, 1\}, \{a, b, c, d\} \times \{0, 1\}) = P \times P$$

is the graph of the self homeomorphism (smooth away from the corners) of the pillowcase

$$(24) \quad c_\delta : P \rightarrow P, \quad c_\delta(\gamma, \theta) = (\gamma, \theta + 2f(\gamma + \pi)).$$

Using the 1-parameter family of perturbations $tf, t \in [0, 1]$ gives an isotopy from the identity $Id : P \rightarrow P$ to $c_\delta : P \rightarrow P$.

Proof. Let a', b', c', d' and μ_E be based loops in $\pi_1(S^2 \times I \setminus (\{a, b, c, d\} \times I \cup E), s)$ so that a', b', c', d' represent the meridians of the punctures in the other boundary component $S^2 \times \{1\}$, and μ_E denotes the meridian to the perturbation curve E . These curves are illustrated in Figure 17, where, for convenience, the four-punctured sphere is identified with a three-punctured disk.

The curves a, b, c, d, μ_E generate $\pi_1(S^2 \times I \setminus (\{a, b, c, d\} \times I \cup E), s)$ and the relations

$$ba = cd, a' = a, b' = b, c' = \mu_E c \bar{\mu}_E, d' = \mu_E d \bar{\mu}_E,$$

hold. The natural longitude λ_E for E is represented by the homotopy class ba .

As explained in [18, Proposition 3.1] (see Proposition 6.1 above), any representation of $\langle a, b, c, d \mid ba = cd \rangle$ taking a, b, c, d to traceless elements is conjugate to one given by

$$(25) \quad a \mapsto \mathbf{i}, b \mapsto e^{\gamma \mathbf{k}} \mathbf{i}, c \mapsto e^{\theta \mathbf{k}} \mathbf{i}, d \mapsto e^{(\theta - \gamma) \mathbf{k}} \mathbf{i},$$




FIGURE 17.

for some $(\gamma, \theta) \in P$. Thus, to any representation $\rho : \pi_1(S^2 \times I \setminus (\{a, b, c, d\} \times I \cup E)) \rightarrow SU(2)$ sending a, b, c, d to traceless elements, one can associate $(\gamma, \theta) \in P$. Then $\rho(\lambda_E) = \rho(ba) = -e^{\gamma \mathbf{k}} = e^{(\gamma + \pi) \mathbf{k}}$.

If $\rho \in R_\delta(S^2 \times I, \{a, b, c, d\} \times I)$, then ρ satisfies the perturbation condition (see Equation (23)):

$$(26) \quad \rho(\mu_E) = e^{f(\gamma + \pi) \mathbf{k}}.$$

Hence

$$(27) \quad \rho(a') = \mathbf{i}, \quad \rho(b') = e^{\gamma \mathbf{k}} \mathbf{i}, \quad \rho(c') = e^{f(\gamma + \pi) \mathbf{k}} e^{\theta \mathbf{k}} \mathbf{i} e^{-f(\gamma + \pi) \mathbf{k}} = e^{\theta + 2f(\gamma + \pi) \mathbf{k}}$$

Conversely, given any $(\gamma, \theta) \in P$ and $e^{\alpha \mathbf{k}}$ there exists a unique traceless representation $\rho : \pi_1(S^2 \times I \setminus (\{a, b, c, d\} \times I \cup E)) \rightarrow SU(2)$ and $\rho(\mu_E) = e^{\alpha \mathbf{k}}$. This satisfies the perturbation condition, and hence $\rho \in R_\delta(S^2 \times I, \{a, b, c, d\} \times I)$ provided $e^{\alpha \mathbf{k}} = e^{f(\gamma + \pi) \mathbf{k}}$.

We have shown that to each $(\gamma, \theta) \in P$ there exists a unique $\rho \in R_\delta(S^2 \times I, \{a, b, c, d\} \times I)$, given by (25), (26), and (27). Moreover the restriction

$$R_\delta(S^2 \times I, \{a, b, c, d\} \times I) \rightarrow R(S^2 \times \{0, 1\}, \{a, b, c, d\} \times \{0, 1\}) = P \times P$$

has image $(\gamma, \theta, \gamma, \theta + 2f(\gamma + \pi))$

This shows that $d = (E, f)$ induces the map $c_\delta(\gamma, \theta) = (\gamma, \theta + 2f(\gamma + \pi))$, as asserted. This map is invertible, with inverse $(\gamma, \theta) \mapsto (\gamma, \theta - 2f(\gamma + \pi))$, and hence is a homeomorphism. \square

We stated Theorem 9.1 for a specific curve e in $S^2 \setminus \{a, b, c, d\}$ but one may conjugate by any diffeomorphism ϕ of the punctured sphere to replace e by $\phi(e)$, generating many more homeomorphisms of the pillowcase. Although not used in the rest of this article, these perturbations will be important in forthcoming work.

Theorem 9.2. *Given any relatively prime pair of integers p, q and $\phi \in \mathcal{X}$, there exists a holonomy perturbation along a single curve in $(S^2 \times I, \{a, b, c, d\} \times I)$ inducing the homeomorphism*

$$c_{p,q,f}(\gamma, \theta) = (\gamma - q\phi(p\gamma + q\theta), \theta + p\phi(p\gamma + q\theta))$$

of the pillowcase. This homeomorphism is Hamiltonian isotopic to the identity.

Proof. Given $(\gamma, \theta) \in \mathbb{R}^2$, Let $\psi(\gamma, \theta) : \pi_1(S^2 \setminus \{a, b, c, d\}) \rightarrow SU(2)$ be the traceless representation of Proposition 6.1.

Let $g : (S^2, \{a, b, c, d\}) \rightarrow (S^2, \{a, b, c, d\})$ be the half-Dehn twist diffeomorphism supported in the hemisphere containing c and d which sends c to d and d to c . (Thus g^2 is the Dehn twist about the curve labeled e in Figure 16.) Choosing a base point near a , the induced automorphism on $\pi_1(S^2 \setminus \{a, b, c, d\})$ is given by

$$g_*(a) = a, g_*(b) = b, g_*(c) = d, g_*(d) = d^{-1}cd.$$

Then $\psi(\gamma, \theta)(g_*(b)) = e^{\gamma \mathbf{k} \mathbf{i}}$ and $\psi(\gamma, \theta)(g_*(c)) = e^{(\theta - \gamma) \mathbf{k} \mathbf{i}}$, i.e.,

$$g^* \psi(\gamma, \theta) = (\gamma, \theta - \gamma).$$

In other words, g induces the linear map on the pillowcase:

$$(28) \quad g^* \begin{pmatrix} \gamma \\ \theta \end{pmatrix} = A_g \begin{pmatrix} \gamma \\ \theta \end{pmatrix}, \quad A_g = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}.$$

Let $h : (S^2, \{a, b, c, d\}) \rightarrow (S^2, \{a, b, c, d\})$ be the diffeomorphism which fixes (a neighborhood of) a , and cyclically permutes b, c, d . This can be chosen to induce the automorphism $\pi_1(S^2 \setminus \{a, b, c, d\})$ is given by

$$h_*(a) = a, h_*(b) = c^{-1}, h_*(c) = d, h_*(d) = (cd)^{-1}b^{-1}(cd)$$

Then $\psi(\gamma, \theta)(h_*(b)) = -e^{\theta \mathbf{k} \mathbf{i}} = e^{(\theta + \pi) \mathbf{k} \mathbf{i}}$ and $\psi(\gamma, \theta)(g_*(c)) = e^{(\theta - \gamma) \mathbf{k} \mathbf{i}}$, i.e., h induces the affine map on the pillowcase:

$$(29) \quad h^* \begin{pmatrix} \gamma \\ \theta \end{pmatrix} = A_h \begin{pmatrix} \gamma \\ \theta \end{pmatrix} + \mathbf{v}, \quad A_h = \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} \pi \\ 0 \end{pmatrix}.$$

The matrices

$$S = A_g A_h^4 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad T = A_h A_g^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

are the standard generators of the modular group. It follows that given any relatively prime pair of integers p, q , there exists a word $w = w(g, h)$ in g and h so that the resulting diffeomorphism $w : (S^2, \{a, b, c, d\}) \rightarrow (S^2, \{a, b, c, d\})$ satisfies

$$(30) \quad w^* \begin{pmatrix} \gamma \\ \theta \end{pmatrix} = A \begin{pmatrix} \gamma \\ \theta \end{pmatrix} + \mathbf{u}, \quad A = \begin{pmatrix} p & q \\ r & s \end{pmatrix}$$

where $ps - qr = 1$ and \mathbf{u} is a vector whose entries are integer multiples of π .

The diffeomorphism w induces a level preserving diffeomorphism

$$w : (S^2 \times I, \{a, b, c, d\} \times I) \rightarrow (S^2 \times I, \{a, b, c, d\} \times I).$$

This diffeomorphism takes perturbed flat connections with respect to the perturbation curve E of Theorem 9.1 to perturbed flat connections with respect to $w(E)$.

To simplify notation, write $\phi(x) = 2f(x + \pi) + u_1$ where $f \in \mathcal{X}$ is the function used in Theorem 9.1, and u_1 is the first component of the vector \mathbf{u} . Note that $\phi \in \mathcal{X}$ if and only if $f \in \mathcal{X}$. Then the self-homeomorphism of the pillowcase given by perturbing along $w(E)$ is the conjugate $(w^*)^{-1} \circ c_\delta \circ w^*$, which we compute

$$\begin{aligned}
((w^*)^{-1} \circ c_\delta \circ w^*) \begin{pmatrix} \gamma \\ \theta \end{pmatrix} &= (w^*)^{-1} \circ c_\delta \left(A \begin{pmatrix} \gamma \\ \theta \end{pmatrix} + \mathbf{u} \right) \\
&= (w^*)^{-1} \left(A \begin{pmatrix} \gamma \\ \theta \end{pmatrix} + \mathbf{u} + \begin{pmatrix} 0 \\ \phi(p\gamma + q\theta + u_1) \end{pmatrix} \right) \\
&= A^{-1} \left(A \begin{pmatrix} \gamma \\ \theta \end{pmatrix} + \mathbf{u} + \phi(p\gamma + q\theta + u_1) \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) - A^{-1} \mathbf{u} \\
&= \begin{pmatrix} \gamma \\ \theta \end{pmatrix} + \phi(p\gamma + q\theta + u_1) \begin{pmatrix} -q \\ p \end{pmatrix}.
\end{aligned}$$

If u_1 is an even multiple of π , then we are done, since these are pillowcase coordinates. If u_1 is an odd multiple of π , replace $\phi(x)$ by $\phi(x + \pi)$; this induces a bijection of \mathcal{X} . \square

10. HOLONOMY PERTURBATIONS TO SMOOTH THE TRACELESS CHARACTER VARIETY OF A 2-TANGLE DECOMPOSITION OF A TORUS KNOT


For the rest of this article we provide a detailed study of the traceless character varieties associated to a certain 2-tangle decomposition of a torus knot $T_{p,q}$. In the present section we establish that $(S^3, T_{p,q})$ admits a 2-tangle decomposition as in Equation 17 which verifies Conjecture 6.3 except possibly for the absence of fishtails.

In the next section we identify $L_1 : R_\pi(Y, T) \rightarrow P$ for a number of $T_{p,q}$ and verify that in all our examples, L_1 is indeed a restricted immersed 1-manifold, and that $H^1(Y, T)$ is either isomorphic to $I^1(S^3, T_{p,q})$, or, in examples where the calculation of $I^1(S^3, T_{p,q})$ is unknown, that the calculation of $H^1(Y, T)$ combined with Conjecture 6.5 is consistent with the conjecture [22] that the ranks of $I^1(S^3, K)$ and knot Heegaard-Floer homology $\widehat{HFK}(K)$ are equal.

We recall the description of the traceless $SU(2)$ character variety of a tangle associated to the (p, q) -torus knot from [18, 17]. Figure 18 illustrates a 3-component link $H_A \cup H_B \cup K$ in S^3 , with the component K intersecting a 3-ball D in a trivial 2-tangle U . Integers r, s satisfying $pr + qs = 1$ are fixed throughout.

Performing $-\frac{s}{p}$ Dehn surgery on the component labeled H_A and $\frac{q}{r}$ Dehn surgery on the component labeled H_B yields S^3 again, and the knot labeled K becomes the (p, q) torus knot. In this S^3 , let Y denote the complement of the illustrated 3-ball D , and T the part of the (p, q) torus knot contained in Y . Precisely, Y is obtained from $S^3 \setminus D$ by performing $-\frac{s}{p}$ and $\frac{q}{r}$ surgery on H_A and H_B , and $T \subset Y$ denotes that part of K which lies in Y . Note that Y is itself diffeomorphic to a 3-ball.

Let P_A and P_B in Y be the cores of the Dehn surgery solid tori which are added after neighborhoods of H_A and H_B are removed. We will perform holonomy perturbations along these curves in Y .

FIGURE 18. The (p, q) torus knot

Generators a, b, c, d, x, y of the fundamental group

$$\pi_1(Y \setminus (T \cup P_A \cup P_B)) = \pi_1(S^3 \setminus (D \cup K \cup H_A \cup H_B))$$

are illustrated.

One computes (see [18]) that

$$\pi_1(Y \setminus (T \cup P_A \cup P_B)) = \langle x, y, a, b, c \mid c = \bar{x}ax, ad\bar{a} = yxb\bar{x}\bar{y}, [y, xb] = 1, [x, d\bar{a}y] = 1 \rangle.$$

The curves

$$A_1 = (xb)^q y^r \text{ and } A_2 = (xb)^{-p} y^s$$

form a longitude-meridian pair for the component P_A . The curves

$$B_1 = (d\bar{a}y)^{-s} x^p \text{ and } B_2 = (d\bar{a}y)^r x^q$$

form a (commuting) longitude-meridian pair for the component P_B . In particular, $\pi_1(Y \setminus T)$ is obtained from $\pi_1(Y \setminus (T \cup P_A \cup P_B))$ by killing A_2 and B_2 .

Working with the presentation of $\pi_1(Y \setminus (T \cup P_A \cup P_B))$ given above, together with the fact that $pr + qs = 1$ yields:

$$A_1^s = (xb)^{qs} y^{rs} = (xb)(xb)^{-pr} y^{rs} = xb((xb)^{-p} y^s)^r = xbA_2^r$$

and so

$$xb = A_1^s A_2^{-r}.$$

Similar calculations give

$$y = A_2^q A_1^p, \quad x = B_2^s B_1^r, \quad d\bar{a}y = B_1^{-q} B_2^p,$$

from which one obtains

$$(31) \quad a = yxb\bar{x}(d\bar{a}y)^{-1} = A_1^{s+p} A_2^{q-r} B_1^{q-r} B_2^{-(s+p)} \text{ and } b = \bar{x}xb = B_1^{-r} B_2^{-s} A_1^s A_2^{-r}.$$

Since $c = \bar{x}ax$ and $d = \bar{a}yxb\bar{x}\bar{y}a$, it follows that the four elements A_1, A_2, B_1, B_2 generate $\pi_1(Y \setminus (T \cup P_A \cup P_B))$. A simple extension of the observation in [18] that $\pi_1(Y \setminus T)$ is free on A_1 and B_1 (they are labeled A and B in that article) shows that $\pi_1(Y \setminus (T \cup P_A \cup P_B))$ is the free product of the free abelian group generated by A_1, A_2 and the free abelian group generated by B_1, B_2 .

We use the perturbation functions $\epsilon_A \sin x$ on P_A and $\epsilon_B \sin x$ on P_B for some $(\epsilon_1, \epsilon_2) \in \mathbb{R}^2$. Recall from Equation (23) that with this choice, perturbed-flat connections modulo gauge are identified with representations $\rho : \pi_1(Y \setminus (T \cup P_A \cup P_B)) \rightarrow SU(2)$ which satisfy the *perturbation conditions*

$$(32) \quad \begin{aligned} \rho(A_2) &= e^{\epsilon_A \sin u Q_A} \text{ if } \rho(A_1) = e^{u Q_A} \text{ for some } Q_A \in C(\mathbf{i}) \\ \rho(B_2) &= e^{\epsilon_B \sin v Q_B} \text{ if } \rho(B_1) = e^{v Q_B} \text{ for some } Q_B \in C(\mathbf{i}). \end{aligned}$$

If $(\epsilon_A, \epsilon_B) = (0, 0)$, then perturbed-flat connections send A_2 and B_2 to $1 \in SU(2)$, hence by the Seifert-Van Kampen theorem correspond exactly to $SU(2)$ representations of $\pi_1(Y \setminus T)$.

As above, we define the *perturbed traceless flat moduli space*

$$R_{\epsilon_A, \epsilon_B}(Y, T) = \{\rho : \pi_1(Y \setminus (T \cup P_A \cup P_B)) \rightarrow SU(2) \mid \rho \text{ traceless, satisfying (32)}\} / \text{conjugation}$$

Theorem 10.1. *There exists a neighborhood $\mathcal{O} \subset \mathbb{R}^2$ of $(0, 0)$ such that for any $(\epsilon_A, \epsilon_B) \in \mathcal{O}$, the space $R_{\epsilon_A, \epsilon_B}(Y, T)$ is a smooth compact 1-manifold with two boundary points and such that the restriction map to the pillowcase $R_{\epsilon_A, \epsilon_B}(Y, T) \rightarrow P$ satisfies the conditions to be a restricted immersed 1-manifold except possibly the absence of fishtails.*

In extensive calculations we have not found any small non-zero ϵ_A, ϵ_B for which $R_{\epsilon_A, \epsilon_B}(Y, T) \rightarrow P$ is *not* a restricted immersed 1-manifold.

The strategy to prove Theorem 10.1 is standard: we form a parameterized moduli space, prove it is a smooth manifold, and apply Sard's theorem to the projection to \mathbb{R}^2 . We start with a *gauge fixing* theorem which identifies $R_{\epsilon_A, \epsilon_B}(Y, T)$ with a subset of the box $[0, \pi] \times [0, \pi] \times [-1, 1]$.

A representation $\rho : \pi_1(Y \setminus (T \cup P_A \cup P_B)) \rightarrow SU(2)$ satisfying the perturbation conditions with respect to (ϵ_A, ϵ_B) is traceless if and only if $\rho(a)$ and $\rho(b)$ are traceless. From Equation (31) this holds if and only if

$$\operatorname{Re}(\rho(A_1^{s+p} A_2^{q-r} B_1^{q-r} B_2^{-(s+p)})) = 0 \text{ and } \operatorname{Re}(\rho(B_1^{-r} B_2^{-s} A_1^s A_2^{-r})) = 0.$$

Assuming that $\rho(A_1) = e^{u Q_A}$ and $\rho(B_1) = e^{v Q_B}$ for some pair of purely imaginary unit quaternions $Q_A, Q_B \in C(\mathbf{i})$, these can be expressed equivalently as

$$\operatorname{Re} \left(e^{((s+p)u + (q-r)\epsilon_A \sin u) Q_A} e^{((q-r)v - (s+p)\epsilon_B \sin v) Q_B} \right) = 0$$

and

$$\operatorname{Re} \left(e^{(-rv - s\epsilon_B \sin v) Q_B} e^{(su - r\epsilon_A \sin u) Q_A} \right) = 0$$

or equivalently (see [18, Proposition 2.1]) as $\Psi(\epsilon_A, \epsilon_B, u, v, \tau) = (0, 0)$, where $\Psi = (\Psi_1, \Psi_2)$ is defined by

$$(33) \quad \begin{aligned} \Psi_1(\epsilon_A, \epsilon_B, u, v, \tau) &= \cos((q-r)v - (s+p)\epsilon_B \sin v) \cos((s+p)u + (q-r)\epsilon_A \sin u) \\ &\quad - \sin((q-r)v - (s+p)\epsilon_B \sin v) \sin((s+p)u + (q-r)\epsilon_A \sin u) \tau \end{aligned}$$

and

$$(34) \quad \begin{aligned} \Psi_2(\epsilon_A, \epsilon_B, u, v, \tau) = & \cos(-rv - s\epsilon_B \sin v) \cos(su - r\epsilon_A \sin u) \\ & - \sin(-rv - s\epsilon_B \sin v) \sin(su - r\epsilon_A \sin u)\tau, \end{aligned}$$

with τ the cosine of the angle between Q_A and Q_B .

Theorem 10.2. *Fix $(\epsilon_A, \epsilon_B) \in \mathbb{R}^2$ and let*

$$W_{\epsilon_A, \epsilon_B} = \{(u, v, \tau) \in [0, \pi] \times [0, \pi] \times [-1, 1] \mid \Psi(\epsilon_A, \epsilon_B, u, v, \tau) = 0\}.$$

Then the assignment

$$A_1 \mapsto e^{ui}, A_2 \mapsto e^{\epsilon_A \sin ui}, B_1 \mapsto e^{ve^{\arccos \tau} \mathbf{k}_i}, B_2 \mapsto e^{\epsilon_B \sin ve^{\arccos \tau} \mathbf{k}_i}$$

defines a surjection

$$f : W_{\epsilon_A, \epsilon_B} \rightarrow R_{\epsilon_A, \epsilon_B}(Y, T)$$

whose fiber over $f(u, v, \tau)$ is the single point $\{(u, v, \tau)\}$ unless $\sin u = 0$ or $\sin v = 0$, in which case the fiber is the arc $\{(u, v)\} \times [-1, 1]$. The map f sends the points of $W_{\epsilon_A, \epsilon_B}$ interior to the box to non-abelian perturbed representations and boundary points to perturbed abelian representations.

The proof of this lemma is identical to [18, Theorem 11.1] (see also [17, Theorem 4.2]). The essential point is that if $\sin u \neq 0 \neq \sin v$ then any representation can be uniquely conjugated so that A_1 is sent to e^{ui} and B_1 is sent to $e^{ve^{\arccos \tau} \mathbf{k}_i}$ for τ the cosine of the angle between this representation's Q_A and Q_B . The perturbation condition then determines where A_2 and B_2 are sent. We leave the details to the reader.

A point $(u, v, \tau) \in W_{\epsilon_A, \epsilon_B}$ which lies on the boundary of the box corresponds to a representation which sends A_1 and A_2 to the center ± 1 if $\sin u = 0$, sends B_1 and B_2 to ± 1 if $\sin v = 0$, and sends A_1, A_2, B_1, B_2 to commuting elements if $|\tau| = 1$. It follows that points in $W_{\epsilon_A, \epsilon_B}$ meeting the boundary of the cube correspond exactly to abelian representations (i.e., representations with abelian image). There are two conjugacy classes of (unperturbed) traceless abelian representations. This property is stable with respect to small perturbations, as shown in Proposition 8.3.

The result needed to prove complete the proof of Theorem 10.1 is the following. It says that the map Ψ of Equations (33), (34) is submersive near non-abelian points of the unperturbed traceless character variety $R(Y, T)$. Hence for generic small (ϵ_A, ϵ_B) , the non-abelian part of $R_{\epsilon_A, \epsilon_B}(Y, T)$ is smooth.

Lemma 10.3. *Suppose that u, v, τ are chosen so that $\Psi(0, 0, u, v, \tau) = 0$, with $\sin u \neq 0, \sin v \neq 0$, and $|\tau| \neq 1$. Then $d\Psi_{(0, 0, u, v, \tau)} : \mathbb{R}^5 \rightarrow \mathbb{R}^2$ is surjective, and hence Ψ is a submersion near $(0, 0, u, v, \tau)$.*

Proof. The proof is essentially a lengthy second-year calculus computation, and we recommend the reader skip it.

Consider first Ψ_2 . To clarify, we adopt the following notation

$$A = \cos(-rv), B = \sin(-rv), C = \cos(su), D = \sin(su).$$

From Equation (34), the partial derivatives of Ψ_2 at the point $(0, 0, u, v, \tau)$ are given by

$$\left. \frac{\partial \Psi_2}{\partial \epsilon_A} \right|_{(0, 0, u, v, \tau)} = A(-D)(-r) \sin u - BC(-r)\tau \sin u = r(AD + BC\tau) \sin u$$

$$\begin{aligned}
\left. \frac{\partial \Psi_2}{\partial \epsilon_B} \right|_{(0,0,u,v,\tau)} &= -B(-s \sin v)C - A(-s \sin v)D\tau = s(BC + AD\tau) \sin v \\
\left. \frac{\partial \Psi_2}{\partial u} \right|_{(0,0,u,v,\tau)} &= A(-D)s - BCs\tau = -s(AD + BC\tau) \\
\left. \frac{\partial \Psi_2}{\partial v} \right|_{(0,0,u,v,\tau)} &= -B(-r)C - A(-r)D\tau = r(BC + AD\tau) \\
\left. \frac{\partial \Psi_2}{\partial \tau} \right|_{(0,0,u,v,\tau)} &= -BD
\end{aligned}$$

Moreover, the equation $\Psi_2(0, 0, u, v, \tau) = 0$ is equivalent to

$$AC - BD\tau = 0.$$

Similarly, adopting the notation

$$E = \cos((q - r)v), F = \sin((q - r)v), G = \cos((s + p)u), H = \sin((s + p)u),$$

we obtain

$$\begin{aligned}
\left. \frac{\partial \Psi_1}{\partial \epsilon_A} \right|_{(0,0,u,v,\tau)} &= -(q - r)(EH + FG\tau) \sin u \\
\left. \frac{\partial \Psi_1}{\partial \epsilon_B} \right|_{(0,0,u,v,\tau)} &= (s + p)(FG + EH\tau) \sin v \\
\left. \frac{\partial \Psi_1}{\partial u} \right|_{(0,0,u,v,\tau)} &= -(s + p)(EH + FG\tau) \\
\left. \frac{\partial \Psi_1}{\partial v} \right|_{(0,0,u,v,\tau)} &= -(q - r)(FG + EH\tau) \\
\left. \frac{\partial \Psi_1}{\partial \tau} \right|_{(0,0,u,v,\tau)} &= -FH
\end{aligned}$$

Moreover, the equation $\Psi_1(0, 0, u, v, \tau) = 0$ is equivalent to

$$EG - FH\tau = 0.$$

Suppose that Ψ were not a submersion at $(0, 0, u, v, \tau)$. Then $d\Psi_1$ and $d\Psi_2$ are linearly dependent.

If $d\Psi_2 = 0$, then $AD + BC\tau = 0$, $BC + AD\tau = 0$ and $BD = 0$. Therefore $AD = AD\tau^2$. Since $|\tau| \neq 1$, this implies $AD = 0$ and $BC = 0$. If $A = \cos(-rv) = 0$, then $B = \sin(-rv) \neq 0$, so $\cos(su) = C = 0$ and hence $\sin(su) = D \neq 0$, contradicting $BD = 0$. But if $A \neq 0$, $\sin(su) = D = 0$ and so $\cos(su) = C \neq 0$ and hence $B = 0$. But this contradicts $AC - BD\tau = 0$. Therefore, $d\Psi_2 \neq 0$. Similarly, $d\Psi_1 \neq 0$.

Since neither $d\Psi_1$ nor $d\Psi_2$ is zero, there exists a non-zero $\alpha \in \mathbb{R}$ so that $\alpha d\Psi_1 = d\Psi_2$. Comparing the first columns, (i.e., $\frac{\partial}{\partial \epsilon_A}$) and using the fact that $\sin u \neq 0$ one sees

$$AD + BC\tau = \alpha \left(\frac{r - q}{r} \right) (EH + FG\tau).$$

Similarly, comparing third columns gives

$$AD + BC\tau = \alpha \left(\frac{s+p}{s} \right) (EH + FG\tau).$$

Since $\frac{(r-q)s}{(s+p)r} = 1 - \frac{1}{r(s+p)} \neq 1$, and $\alpha \neq 0$, this implies that $EH + FG\tau = 0$, and hence also $AD + BC\tau = 0$. Comparing the second and fourth columns and applying the same argument yields $FG + EH\tau = 0$ and $AD + BC\tau = 0$.

Then $FG = -EH\tau = FG\tau^2$. Since $|\tau| \neq 1$, $FG = 0 = EH$ and, similarly, $AD = BC = 0$. Since neither $d\Psi_1$ nor $d\Psi_2$ is zero, $BD \neq 0$ and $FH \neq 0$. Thus $A = C = 0$ and $G = E = 0$.

Recalling their definitions, this says that

$$(35) \quad \cos(-rv) = \cos(su) = \cos((q-r)v) = \cos((s+p)u) = 0.$$

Hence there exist odd integers k, ℓ so that $-rv = k\frac{\pi}{2}$ and $(q-r)v = \ell\frac{\pi}{2}$. Thus $(q-r)k = -r\ell$. Since r and $q-r$ are relatively prime, there exist odd integers m, n so that $k = rm$ and $\ell = (q-r)n$. Thus $v = \frac{k}{-r}\frac{\pi}{2} = -m\frac{\pi}{2}$. Similarly, u is an odd multiple of $\frac{\pi}{2}$. Equation (35) then implies that $r, s, q-r, s+p$ are all odd, but then p, q are both even, contradicts the fact that p and q are relatively prime. Thus the assumption that Ψ is not a submersion leads to a contradiction.

Hence Ψ is a submersion at $(0, 0, u, v, \tau)$, and so also near $(0, 0, u, v, \tau)$. \square

Proof of Theorem 10.1. Recall that $W_{0,0}$ denotes the preimage in the box $[0, \pi] \times [0, \pi] \times [-1, 1]$ of 0 by the map $(u, v, \tau) \mapsto \Psi(0, 0, u, v, \tau)$. Let V be the intersection of a small open neighborhood of $W_{0,0}$ with the interior of the box.

Lemma 10.3 implies that (after perhaps choosing a smaller V), there is a neighborhood \mathcal{O} of 0 in \mathbb{R}^2 so that $\Psi : \mathbb{R}^2 \times [0, \pi] \times [0, \pi] \times [-1, 1] \rightarrow \mathbb{R}^2$ restricts to a submersion on $\mathcal{O} \times V$. The *parameterized moduli space* $P := \mathcal{O} \times V \cap \Psi^{-1}(0)$ is a smooth submanifold of $\mathcal{O} \times V$. By Sard's theorem, there exist regular values (ϵ_A, ϵ_B) of the composite $P \subset \mathcal{O} \times V \xrightarrow{\pi \circ \Psi} \mathcal{O}$ arbitrarily close to 0. Its preimage in the interior of the box is identified with the non-abelian part of $R_{\epsilon_A, \epsilon_B}(Y, T)$ by Theorem 10.2. The structure near the two abelian representations was identified in Theorem 8.4. \square

11. CALCULATIONS FOR TORUS KNOTS

In this section we carry out calculations of $C^{\natural}(L_0, L_1)$ for some torus knots, including examples with non-trivial differentials. In what follows, we continue to use the description of the torus knot $K = T_{p,q}$ illustrated in Figure 18, where we perform $-\frac{s}{p}$ Dehn surgery on the component labeled H_A and $\frac{q}{r}$ surgery on the component labeled H_B . Figure 18 illustrates a decomposition

$$(S^3, T_{p,q}) = (Y, T) \cup_{(S^2, \{a,b,c,d\})} (D, U).$$

Recall that this decomposition depends on the choice of integers r, s satisfying $pr + qs = 1$. Different choices of r, s lead to different pairs $R(Y, T)$ and restriction maps L_1 .

The identification of the spaces $R_{\pi}(Y, T)$ and their image in the pillowcase was done using a computer algebra package.

Recall that Kronheimer-Mrowka prove ([23, 20], also Lim [26]) that the rank of the reduced singular instanton homology $I^\natural(S^3, K)$ is bounded below by the sum of the absolute value of coefficients of the Alexander polynomial, which we denote by $|\Delta_K|$.

It is conjectured that the reduced instanton homology of a torus knot K has rank *equal* to $|\Delta_K|$. This is a special case of a more sweeping conjecture which relates singular instanton homology of a knot and its Heegaard-Floer homology.

If Conjecture 6.5 is true, then the rank of $H^\natural(Y, T, \pi)$ must be at least as large as $|\Delta_K|$ for a tangle decomposition of a torus knot K , and if in addition the conjecture of the previous paragraph is true, then $\text{rank}(H^\natural(Y, T, \pi)) = |\Delta_K|$. The reader can verify that in all the examples given below, the rank of $H^\natural(Y, T, \pi)$ equals $|\Delta_K|$.

In the following calculations, we take $L_0 = L_0^{\epsilon, 0}$ for a small $\epsilon > 0$. We calculate Maslov indices relative to the slope 1 line field ℓ_1 , making use of Equation (16) to simplify grading calculations. We make frequent use of the calculus described in Section 5.

11.1. The (5, 11) torus knot. Consider the tangle associated to the (5, 11) torus knot, corresponding to the choice $(5, 11, 9, -4)$. The *unperturbed* traceless character variety $R(Y, T)$ is a restricted immersed 1-manifold.

In fact, $R(Y, T)$ is a union of an arc R_0 and four circles R_1, R_2, R_3, R_4 . The arc R_0 embeds linearly with slope 2. Two of the circles, say R_1, R_2 are vertically monotonic with vertical degree 2. The remaining circles R_3, R_4 each map precisely in the way illustrated in the example of Figures 1, 6, 5, and 8.

Since the signature of the (5, 11) torus knot is $-24 \equiv 0 \pmod{4}$, R_0 contributes $(1, 0, 0, 0)$ to $H^\natural(Y, T)$ by our absolute grading convention. Proposition 5.3 implies that the two vertically monotonic circles R_1, R_2 contribute $(1, 1, 1, 1)$ each to $H^\natural(Y, T)$.

The contributions of R_3 and R_4 to $C(L_0, L_1)$ and $H^\natural(Y, T)$ were computed in detail in Section 3.9; it was shown that each contributes a $(2, 2, 2, 2)$ summand to $C(L_0, L_1)$, each has two bigons contributing to the differential, and each contributes $(1, 1, 1, 1)$ to the homology $H^\natural(Y, T)$. Thus the differential $\partial : C(L_0, L_1) \rightarrow C(L_0, L_1)$ has rank 8 and

$$H^\natural(Y, T) = (1, 0, 0, 0) \oplus_{i=1}^4 (1, 1, 1, 1) = (5, 4, 4, 4).$$

In particular, the rank of $H^\natural(Y, T)$ is 17, which equals $|\Delta_K|$. The calculation of $I^\natural(S^3, T_{5,11})$ is unknown, but Conjecture 6.5 would imply that the Kronheimer-Mrowka lower bound is attained for $T_{5,11}$.

11.2. The (3, 7) torus knot. Taking the decomposition of the (3, 7) torus knot corresponding to the choice $(p, q, r, s) = (3, 7, 5, -2)$, it is shown in [17] that the space $R(Y, T)$ is the disjoint union of an arc R_0 (consisting of binary dihedral representations) and two circles R_1, R_2 . In particular, $L_1 : R(Y, T) \rightarrow P$ is a restricted immersed 1-manifold.

The restriction $L_1|_{R_0} : R_0 \rightarrow P$ is the linear embedding of slope 2 ([17, Theorem 4.1]), given by $[0, \pi] \ni t \mapsto (t, 2t) \in P$. The restrictions of L_1 to R_1 and R_2 have the same image, and each is a vertically monotonic circle of vertical degree 2.

The line segment of slope 2 has a unique intersection point with L_0 , namely the point r_+^ϵ . The signature of $T_{3,7}$ is $-8 \equiv 0 \pmod{4}$ and so R_0 contributes $(1, 0, 0, 0)$ to $H^\natural(Y, T)$.

Since R_1 and R_2 are vertically monotonic with vertical degree 2, Proposition 5.3 implies that R_1 and R_2 each contribute a summand $(1, 1, 1, 1)$ to $C(L_0, L_1)$ and the differential is

zero in these summands. Hence

$$H^\natural(Y, T) = (1, 0, 0, 0) \oplus_{i=1}^2 (1, 1, 1, 1) = (3, 2, 2, 2).$$

This agrees with the calculation of the reduced instanton homology of the $(3, 7)$ torus knot $I^\natural(S^3, T_{3,7})$ (see [18]), as well as the calculation of the $(\mathbb{Z}/4)$ graded) reduced Khovanov homology $Kh^{red}(T_{3,7}^m)$.

11.3. The $(5, 7)$ torus knot. It is established in [17] that taking $(p, q, r, s) = (5, 7, 3, -2)$, $R(Y, T)$ is smooth, and has two components, an arc R_0 and a circle R_1 . The restriction of L_1 to R_0 is linear with slope 2. The restriction of L_1 to the circle is a 2-1 cover onto its image which winds four times vertically around P^* . In particular, $L_1 : R_1 \rightarrow P$ is vertically monotonic with vertical degree $d = 8$.

The arc component R_0 has a unique intersection point with L_0 , namely the point r_+^ϵ . The signature of the $(5, 7)$ torus knot is $-16 \equiv 0 \pmod{4}$, and hence R_0 contributes $(1, 0, 0, 0)$ to $C(L_0, L_1)$ and $H^\natural(Y, T)$. Since R_1 is vertically monotonic with vertical degree 8, Proposition 5.3 implies that R_1 contributes $(4, 4, 4, 4)$ to $C(L_0, L_1)$. Moreover, all differentials are zero in this summand. Thus we conclude that all differential vanish and

$$H^\natural(Y, T) = (5, 4, 4, 4).$$

This agrees with the calculation of the reduced instanton homology $I^\natural(S^3, T_{5,7})$. Moreover, the reduced Khovanov homology $Kh^{red}(T_{5,7}^m)$ equals $(8, 6, 7, 8)$, which has strictly larger rank, corresponding to the fact that there are non-trivial higher differentials in the Kronheimer-Mrowka spectral sequence from Khovanov to instanton homology [23].

11.4. The $(5, 12)$ torus knot. Taking $(p, q, r, s) = (5, 12, 5, -2)$, it is shown in [17] that $R(Y, T) \rightarrow P$ is a restricted immersed 1-manifold composed of one arc R_0 mapping with slope 6, and two vertically monotonic circles R_1, R_2 each of vertical degree 6.


Proposition 5.3 shows that the components R_1 and R_2 each contribute $(3, 3, 3, 3)$ to $C(L_0, L_1)$ and $H^\natural(Y, T)$. For the component R_0 , we calculate in exactly the same manner as was done for 2-bridge knots. There are 5 representations, $r_+^\epsilon, x_1^\pm, y_1^\pm$, and $gr(r_+^\epsilon, x_1^\pm) = 2$, $gr(x_1^\pm, x_2^\pm) = 2$, and $gr(x_i^\pm, x_i^\mp) = 1$. The signature of $K_{5,12}$ is -28 , and hence $gr(r_+^\epsilon) = 0 \pmod{4}$. Since R_0 maps linearly with slope 6 to the pillowcase, there can be no bigons, and thus all differentials are zero. Therefore, the component R_0 contributes $(2, 1, 1, 1)$ to $H^\natural(Y, T)$, generated by $\{r_+^\epsilon, x_2^\pm\}, \{x_1^\pm\}, \{x_1^\pm\}, \{x_2^\pm\}$ respectively. We conclude that

$$H^\natural(Y, T) = (2, 1, 1, 1) \oplus_{i=1}^2 (3, 3, 3, 3) = (8, 7, 7, 7).$$

This agrees with the calculation of $I^\natural(S^3, T_{5,12})$ in [18], and is smaller than the reduced Khovanov homology $Kh^{red}(T_{5,12}^m) = (20, 19, 19, 19)$.

11.5. The $(5, 17)$ torus knot. Taking $(5, 17, 7, -2)$, it is shown in [17] that $R(Y, T) \rightarrow P$ is a restricted immersed 1-manifold, the union of an arc R_0 mapping linearly with slope 2 and three circles R_1, R_2, R_3 are vertically monotonic with vertical degrees 6, 6 and 8. The signature equals $-40 \equiv 0 \pmod{4}$, and so the single intersection r_+^ϵ on the arc R_0 contributes $(1, 0, 0, 0)$ to $C(L_0, L_1)$ and $H^\natural(Y, T)$. Proposition 5.3 implies that the circles R_1, R_2, R_3 contribute $(3, 3, 3, 3)$, $(3, 3, 3, 3)$ and $(4, 4, 4, 4)$ respectively to $C(L_0, L_1)$ and $H^\natural(Y, T)$. Hence

$$H^\natural(Y, T) = (11, 10, 10, 10),$$

FIGURE 19. $R(Y, T)$ and its image in the pillowcase for the $(3, 4)$ torus knot.

which agrees with the calculation of the instanton homology $I^\sharp(S^3, T_{5,17})$ in [18]. Two of the circles (the degree 6 circles) have the same image and the degree 8 circle double covers its image.



11.6. The $(3, 4)$ torus knot. We next give an example of a tangle decomposition so that the unperturbed traceless character variety $R(Y, T)$ is not smooth. After smoothing it using a holonomy perturbation, $R_\pi(Y, T)$ becomes a restricted immersed 1-manifold and one finds a non-trivial differential in the summand corresponding to the arc component.

Take $(p, q, r, s) = (3, 4, 3, -2)$, it was shown in [18, Proposition 11.4] that $R(Y, T)$ is a singular space, obtained from 3 arcs $I_0 \cong [0, \pi]$, $I_\pm \cong [\frac{\pi}{6}, \frac{5\pi}{6}]$ by identifying the endpoints of I_+ and I_- to I_0 at $\frac{\pi}{6}$ and $\frac{5\pi}{6}$ to form a singular variety homeomorphic to the letter ϕ . The restriction map to the pillowcase takes the arc I_0 (consisting of binary dihedral representations) to the arc of slope -2 via $[0, \pi] \ni t \mapsto (\pi - t, -2t)$ and takes each of the two arcs I_\pm to linear arcs of slope 4. The space $R(Y, T)$ and its image in the pillowcase is illustrated in Figure 19.

Applying Theorem 10.1, we can find an arbitrarily small perturbation so that $R_{\epsilon_A, \epsilon_B}(Y, T)$ is smooth. A lengthy calculation (or using a computer algebra package) reveals that $R_{\epsilon_A, 0}(Y, T)$ is the union of an arc and a circle for any small non-zero ϵ_A , as illustrated in Figure 20.

The image of the perturbed character variety, a restricted immersed 1-manifold composed of one arc and one circle, is illustrated in Figure 21, along with the image of L_0 . The arc component R_0 intersects L_0 in three points, r_+^ϵ, x_1^+ and x_1^- . The signature of the $(3, 4)$ torus knot is $-6 \equiv 2 \pmod{4}$, and so $gr(r_+^\epsilon) = 2$. The relative gradings are $gr(x_1^+, x_1^-) = 1$ and $gr(x_1^-, r_+^\epsilon) = 1$. Hence $gr(x_1^-) = 3$ and $gr(x_1^+) = 0$.

A bigon from x_1^- to r_+^ϵ is indicated in in Figure 21. This is the only bigon and gives the non-zero differential $\partial x_1^- = r_+^\epsilon$. Thus R_0 contributes $(1, 0, 0, 0)$ to $H^\sharp(Y, T, \pi)$. The circle component R_1 is vertically monotonic of vertical degree 2 and hence contributes $(1, 1, 1, 1)$

FIGURE 20. $R(Y, T)$ resolving to $R_{\epsilon_A, 0}(Y, T)$ for the (3, 4) torus knot.FIGURE 21. The image of $R_{\epsilon_A, 0}(Y, T)$ in the pillowcase for the (3, 4) torus knot. The bigon giving a non-trivial differential is shaded.

to $C(L_0, L_1)$ and $H^\natural(Y, T)$. Hence

$$H^\natural(Y, T, \pi) = (2, 1, 1, 1),$$

which is isomorphic to the reduced instanton homology $I^\natural(S^3, T_{3,4})$, as well as the reduced Khovanov homology $Kh^{red}(T_{3,4}^m)$.

11.7. The (3, 5) torus knot. Taking $(p, q, r, s) = (3, 5, 2, -1)$, the space $R(Y, T)$ is again a (singular) ϕ curve, made up of an arc $I_0 = [0, \pi]$ of traceless binary dihedral representations which maps to the bottom edge of the pillowcase with slope 0: $[0, \pi] \ni t \mapsto (t, 0)$ and two arcs $I_\pm = [\frac{\pi}{6}, \frac{5\pi}{6}]$ whose interiors consist of non-binary dihedral representations and which map to the pillowcase linearly with slope 6: $I_\pm = [\frac{\pi}{6}, \frac{5\pi}{6}] \ni t \mapsto (t, 6(t - \frac{\pi}{6}))$.

However, the singularities resolve differently than in the case of the (3, 4) torus knot which was illustrated in Figure 20: the perturbed variety $R_{\epsilon_A, \epsilon_B}(Y, T)$ is a single arc R_0 . The image of this arc in the pillowcase is illustrated in Figure 22.




FIGURE 22. The image of $R_{\epsilon_A, \epsilon_B}(Y, T)$ and L_0 in the pillowcase for the $(3, 5)$ torus knot. The bigon giving a non-trivial differential is shaded.

One can easily see a bigon from x_1^- to r_+^ϵ . This is the only bigon, so that $\partial x_1^- = r_+^\epsilon$ is the only non-zero differential, and $gr(x_1^-, r_+^\epsilon) = 1$. The signature of the $(3, 5)$ torus knot is -8 , so that $gr(r_+^\epsilon) = 0$. One computes $gr(x_1^-, x_2^-) = gr(x_2^-, x_3^-) = gr(x_3^-, x_4^-) = 2$. Together with $gr(x_\ell^+, x_\ell^-) = 1$, this gives $C(L_0, L_1) = (3, 2, 2, 2)$ and

$$H^\natural(Y, T, \pi) = (2, 1, 2, 2).$$

Once again, this agrees as an absolutely $\mathbb{Z}/4$ graded group, with $I^\natural(S^3, T_{3,5})$ (and $Kh^{red}(T_{3,5}^m)$).

11.8. The $(4, 5)$ torus knot. This example is interesting in the context of instanton homology, as it was shown by Kronheimer-Mrowka [24] that there is a non-trivial higher differential in their spectral sequence from Khovanov homology to instanton homology.

The description of $R(Y, T)$ for the $(4, 5)$ torus knot, corresponding to the tangle decomposition associated to $(p, q, r, s) = (4, 5, 4, -3)$, is analyzed in detail in [17, Section 4]. We refer the reader to that article, where it is shown that $R(Y, T)$ is again a ϕ curve, and its restriction map to P is illustrated, along with its nine intersection points with L_0 generating $C(L_0, L_1)$ and the reduced instanton complex.

A computer-aided calculation shows that $R_{\epsilon_A, 0}(Y, T)$ is the union of an arc R_0 and a circle R_1 and that the restriction to the pillowcase is a restricted immersed 1-manifold. The circle R_1 is vertically monotonic of vertical degree 2. Hence R_1 contributes $(1, 1, 1, 1)$ to $C(L_0, L_1)$ and $H^\natural(Y, T)$.




FIGURE 23. The image of the immersed arc $R_0 \subset R_{\epsilon_A, \epsilon_B}(Y, T)$ and L_0 in the pillowcase for the $(4,5)$ torus knot. The bigon giving a non-trivial differential is shaded.

The image of the arc $R_0 \subset R_{\epsilon_A, 0}(Y, T)$ in the pillowcase is illustrated in Figure 23. There is only one bigon and, in contrast to the examples of the $(3,5)$ and $(4,5)$ torus knots given above, the non-trivial differential does not involve the canonical generator r_+^ϵ . The signature of the $(4,5)$ torus knot is -8 , so that $gr(r_+^\epsilon) = 0$.

One computes that R_0 contributes $(2, 1, 1, 1)$ to $C(L_0, L_1)$. This uses the observation that R_0 has two slope 1 tangencies. The differential takes a generator in grading 1 to a generator in grading 0. Hence R_0 contributes $(1, 0, 1, 1)$ to $H^\natural(Y, T, \pi)$, so that

$$H^\natural(Y, T, \pi) = (2, 1, 2, 2).$$

Once again, this is isomorphic as a $\mathbb{Z}/4$ graded group to $I^\natural(S^3, T_{4,5})$, computed in [24].

It is worth contrasting this calculation with the one Kronheimer-Mrowka give of the instanton homology $I^\natural(S^3, K)$. They start with the count of the nine generators and their relative grading to get the relatively graded chain complex with ranks (up to cyclic reordering) $(3, 2, 2, 2)$. They then compare this to $Kh^{red}(T_{4,5}) = (2, 1, 3, 3)$ to conclude, from the incompatibility of gradings, that there must be a non-trivial differential. A further non-trivial argument identifies this differential. By contrast, the differential is manifest in our pictures. Of course, Conjecture 6.5 may be false, and so one should be cautious in drawing conclusions.




FIGURE 24. $R(Y, T)$ and its smoothing $R_{\epsilon_A, \epsilon_B}(Y, T) = R_0 \cup R_1 \cup R_2$ for the $(4, 7)$ torus knot.




FIGURE 25. The image of the immersed arc $R_0 \subset R_{\epsilon_A, \epsilon_B}(Y, T)$ in the pillowcase for the $(4, 7)$ torus knot. The two bigons are shaded.

11.9. The $(4, 7)$ torus knot. We take the tangle decomposition of the $(4, 7)$ torus knot determined by the choice $(p, q, r, s) = (4, 7, 2, -1)$. The singular variety $R(Y, T)$ and its image in the pillowcase is illustrated in [17, Figure 9]. The smoothed variety $R_{\epsilon_A, \epsilon_B}(Y, T)$ is the union of an arc R_0 and two circles R_1, R_2 . The map to the pillowcase is a restricted immersed 1-manifold.

The two circles each map to vertically monotonic circles of vertical degree 2; each contributes $(1, 1, 1, 1)$ to $H^{\natural}(Y, T, \pi)$. The restriction map of the arc R_0 to the pillowcase is illustrated in Figure 25. There are seven intersection points, and two bigons are shaded. Notice also the four points where the tangent line of $L_1(R_0)$ is tangent to the slope 1 line field ℓ_1 . The $(4, 7)$ torus knot has signature $-14 \equiv 2 \pmod{4}$, so that $gr(r_+^{\epsilon}) = 2$. From this one computes that the contribution of R_0 to $C(L_0, L_1)$ is $(2, 1, 2, 2)$ and to $H^{\natural}(Y, T, \pi)$

is $(1, 0, 1, 1)$. Therefore,

$$H^{\natural}(Y, T, \pi) = (3, 2, 3, 3).$$

In particular, the rank is 11, equal to the sum of the absolute value of the coefficients of the Alexander polynomial of $T_{4,7}$.

The calculation of $I^{\natural}(S^3, T_{4,7})$ is unknown to us. Notice that $C(L_0, L_1) = (4, 3, 4, 4)$ (with one non-trivial differential from grading 1 to 0 and the other from grading 3 to 2). This is consistent with the relative gradings computed for the instanton chain complex in [18]. In that article we computed gradings using a spectral flow splitting formula approach (suggested in [23]), based, not on a tangle decomposition as in the current article, but rather on the decomposition of the form $(S^3, K) = (S^3 \setminus N(K), \phi) \cup_{T^2} (N(K), K)$. This yielded $CI^{\natural}(S^3, T_{4,7}) = (1, 0, 0, 0)_a \oplus (4, 4, 3, 3)_b$, where the subscripts denote a possible cyclic re-ordering. We conjecture in that article that $a = \sigma(K)$ and $b = 3$, which implies that $CI^{\natural}(S^3, T_{4,7}) = (0, 0, 1, 0) \oplus (4, 3, 3, 4) = (4, 3, 4, 4)$. Thus, although we do not know the instanton homology, we do see that the generators of the instanton chain complex occur in the same gradings as for $C(L_0, L_1)$.

11.10. Changing ϵ to cancel bigons. Consider the effect of varying ϵ in the definition for $L_0^{\epsilon,0} : S^1 \rightarrow P^*$ in Definition 13 and illustrated in Figure 7.

For very small $\epsilon > 0$, there are $2n + 1$ intersection points of $L_1(R(Y, T))$ and $L_0^{\epsilon}(R(D, U))$, where n corresponds to the number of intersections of $L_1(R(Y, T))$ with the diagonal arc Δ , or equivalently (see [18]) n equals the number of conjugacy classes of non-abelian traceless representations of the corresponding knot complement. This is doubled to account for the fact that $L_0^{\epsilon,0}$ is a figure 8 close to Δ , and the extra intersection point corresponds to the the perturbation of the unique abelian traceless representation which is mapped to the corner. An illustration of this is given in Figure 15, where one sees $10 = 2 \cdot 5$ points, labeled x_i^{\pm} , corresponding to the five intersection points with the interior of Δ , and one extra point r_+^{ϵ} which converges to the bottom left corner as $\epsilon \rightarrow 0$.

Increasing the parameter ϵ in the holonomy perturbation function makes the figure 8 $L_0^{\epsilon,0}$ wider (see [18]). In some circumstances, the regular homotopy of $L_0^{\epsilon,0}$ obtained by increasing ϵ can be used to cancel pairs of intersection points, and hence reduce the rank $C(L_0, L_1)$.

For example, for the $(3, 5)$ torus knot, a bigon is illustrated in Figure 22. As ϵ increases, the pair of intersection points r_+^{ϵ} and x_1^- get closer together and eventually cancel. Explicitly, when $\epsilon = 0.2$, $\epsilon_A = 0.1$, and $\epsilon_B = 0$, $L_1(R_{\epsilon_A,0}(Y, T))$ intersects L_0 in 7 points. This is the minimum possible by the lower bound given by the sum of absolute value of the coefficients of the Alexander polynomial $t^8 - t^7 + t^5 - t^4 + t^3 - t + 1$ of the $(3, 5)$ torus knot, since, by Proposition 6.4, the rank of $C(L_0^{\epsilon,g}, L_1)$ cannot be smaller than the rank of the instanton homology.

The same method works for the tangle decomposition of the $(3, 4)$ torus knot with $R(Y, T)$ illustrated in Figure 19; increasing ϵ to 0.8 removes the two generators spanned by a bigon. This shows that a suitably perturbed Chern-Simons function on the configuration space of the $(3, 4)$ torus knot is perfect.

We summarize:

Proposition 11.1. *There exists a holonomy perturbations $CS + h$ of the Chern-Simons function on the orbit space of singular connections on $(S^3, T_{3,4})$ and on $(S^3, T_{3,5})$ so that $CS + h$ is perfect, and hence all differentials in the singular instanton complex vanish. \square*

In general, simply increasing ϵ does not eliminate every pair of generators spanned by a bigon. For example, in the 2-tangle decomposition of the $(4, 5)$ torus knot, increasing ϵ increases the number of intersection points of L_0^ϵ with L_1 . For the $(4, 7)$ torus knot, one pair (r_+^ϵ and x_3^- in Figure 25) can be eliminated by increasing ϵ but the second pair (x_2^+ and x_1^-) cannot.

Theorem 4.1 implies that $H^\natural(L_0, L_1)$ is unchanged by a homotopy of L_0 , and, in particular, bigons can be used as guides to regularly homotop away pairs of intersection points. For example, in the case of the $(4, 7)$ torus knot, one can easily find a curve L'_0 homotopic to L_0^ϵ so that all differentials in the resulting complex are zero. It is not clear whether such an L'_0 can be found only using holonomy perturbations.

12. LOOSE ENDS

Several problems remain to be settled before our approach can be considered as producing a functioning invariant of knots. Conjectures 6.3 and 6.5 need to be further investigated.

An important first problem is to determine the extent to which $H^\natural(Y, T, \pi)$ depends on the perturbation π . Given two perturbations π, π' for which $R_\pi(Y, T) \rightarrow P$ and $R_{\pi'}(Y, T) \rightarrow P$ are both restricted immersed 1-manifolds, they are not necessarily related by a regular homotopy, but rather by a *Legendrian cobordism* [14]. For example, in the calculations with torus knots described in Section 11, we used the perturbation corresponding to $\epsilon_A > 0, \epsilon_B = 0$ with ϵ_A small to smooth $R(Y, T)$. Typically, using $\epsilon_A < 0, \epsilon_B = 0$ resolves the normal crossing singularities along the arc of binary dihedrals in the opposite way. For these examples, the resulting $H^\natural(Y, T, \pi)$ is unchanged by reversing the sign of ϵ_A . But in general, Legendrian cobordisms need not preserve Lagrangian-Floer homology.

A closely related question concerns the existence of fishtails, which obstruct $\partial^2 = 0$. We would like to know that there are no fishtails for small perturbations π .

A third question concerns the relationship of $H^\natural(Y, T, \pi)$ to reduced Khovanov homology, a question already solved for Heegaard-Floer theory in [28] and for singular instanton homology in [23]. In forthcoming work we explore this question, extending the definition of $H^\natural(Y, T)$ to include links, and we have established a skein exact triangle for $H^\natural(Y, T, \pi)$. We expect this to lead to a spectral sequence from Khovanov homology to $H^\natural(Y, T, \pi)$ and to an approach to prove Conjecture 6.3.

A fourth question concerns the promoting of the constructions of this article to n -tangle decompositions of knots and links. Some related work includes [19], which studies decompositions of a knot into two trivial n -tangles. The symplectic variety corresponding to the pillowcase in this setting is no longer 2-dimensional, making it much more difficult to understand and compute with.

In a different direction, the rich collections of isotopies of the pillowcase described in Theorems 9.1 and 9.2 are induced by holonomy perturbations, which also induce analytically appropriate perturbations of the Chern-Simons functional for the construction of instanton homology. These should prove useful in isotoping L_0^ϵ to reduce the number of generators of the instanton chain complex.

REFERENCES

- [1] M. Abouzaid, *On the Fukaya categories of higher genus surfaces*. Adv. Math. 217 (2008), no. 3, 1192–1235.
- [2] M. Atiyah, *New invariants of 3- and 4-dimensional manifolds*. “The mathematical heritage of Hermann Weyl (Durham, NC, 1987)”. Proc. Sympos. Pur Math. 48. Amer. Math. Soc., Providence, RI, (1988). 285–299.
- [3] M. Atiyah, *Floer homology*. “The Floer memorial volume.” Progr. Math. 133. Birkhäuser, Basel. (1995). 105–108.
- [4] D. Auroux, *A beginner’s introduction to Fukaya categories*. <http://arxiv.org/abs/1301.7056>
- [5] H. Boden, C. Herald, P. Kirk, and E. Klassen, *Gauge theoretic invariants of Dehn surgeries on knots*. Geom. Topol. 5 (2001), 143–226.
- [6] P. Buser, “Geometry and spectra of compact Riemann surfaces.” Progress in Mathematics, 106. Birkhäuser Boston, Inc., Boston, MA, (1992). xiv+454 pp.
- [7] S. Cappell, R. Lee, and E. Miller, *On the Maslov index*. Comm. Pure Appl. Math. 47 (1994), no. 2, 121–186.
- [8] D. Duncan, *Higher-rank instanton cohomology and the quilted Atiyah-Floer conjecture* <http://arxiv.org/abs/1311.5609>
- [9] A.M. Daniel and P. Kirk, *A general splitting formula for the spectral flow*. With an appendix by K. P. Wojciechowski. Michigan Math. J. 46 (1999), no. 3, 589–617.
- [10] S. Donaldson, *The orientation of Yang-Mills moduli spaces and 4-manifold topology*. J. Differential Geom. 26 (1987), no. 3, 397–428.
- [11] S. Donaldson, “Floer homology groups in Yang-Mills theory.” With the assistance of M. Furuta and D. Kotschick. Cambridge Tracts in Mathematics, 147. Cambridge University Press, Cambridge, 2002. viii+236 pp.
- [12] S. Dostoglou and D. Salamon, *Self-dual instantons and holomorphic curves*. Ann. of Math. (2) 139 (1994), no. 3, 581–640.
- [13] D.B.A. Epstein, *Curves on 2-manifolds and isotopies*. Acta Math. 115 1966 83–107.
- [14] C. Herald, *Legendrian cobordism and Chern-Simons theory on 3-manifolds with boundary*. Comm. Anal. Geom. 2 (1994), no. 3, 337–413.
- [15] C. Herald, *Flat connections, the Alexander invariant, and Casson’s invariant*. Comm. Anal. Geom. 5 (1997), no. 1, 93–120.
- [16] A. Floer, *Morse theory for Lagrangian intersections*. J. Differential Geometry 28 (1988), 513–547.
- [17] Y. Fukumoto, P. Kirk, and J. Pinzon-Caicédo, *Traceless $SU(2)$ representations of 2-stranded tangles*. <http://arxiv.org/abs/1305.6042>
- [18] C. Herald, M. Hedden, and P. Kirk, *The pillowcase and perturbations of traceless representations of knot groups*. Geom. Topol. 18 (2014), no. 1, 211–287.
- [19] Magnus Jacobsson, and Ryszard L. Rubinsztein, *Symplectic topology of $SU(2)$ -representation varieties and link homology, I: Symplectic braid action and the first Chern class*. <http://arxiv.org/abs/0806.2902>.
- [20] P. Kronheimer and T. Mrowka, *Instanton Floer homology and the Alexander polynomial*. Algebr. Geom. Topol. 10 (2010), no. 3, 1715–1738.
- [21] P. Kronheimer and T. Mrowka, *Knot homology groups from instantons*. J. Topol. 4 (2011), no. 4, 835–918.
- [22] P. Kronheimer and T. Mrowka, *Knots, sutures, and excision*. J. Differential Geom. 84 (2010), no. 2, 301–364.
- [23] P. Kronheimer and T. Mrowka, *Khovanov homology is an unknot-detector*. Publ. Math. Inst. Hautes Études Sci. No. 113 (2011), 97–208.
- [24] P. Kronheimer and T. Mrowka, *Filtrations on instanton homology*. <http://arxiv.org/abs/1110.1290>.
- [25] L. Nicolaescu, *The Maslov index, the spectral flow, and splittings of manifolds*, Duke Math. J. 80 (1995) 485–533.
- [26] Y. Lim, *Instanton homology and the Alexander polynomial*. Proc. Amer. Math. Soc. 138 (2010), no. 10, 3759–3768.

- [27] A. Lobb and R. Zentner, *On spectral sequences from Khovanov homology*. <http://arxiv.org/abs/1310.7909>.
- [28] P. Ozsváth, and Z. Szabó, *On the Heegaard Floer homology of branched double-covers*. Adv. Math. 194 (2005), no. 1, 1–33.
- [29] V. de Silva, J. Robbin and D. Salamon, *Combinatorial Floer homology*. Memoirs of the American Mathematical Society Volume 230, Number 1080.
- [30] N. Saveliev, *Floer homology of Brieskorn homology spheres*. J. Differential Geom. 53 (1999), no. 1, 15–87.
- [31] P. Seidel, *Graded Lagrangian submanifolds*. Bull. de la S.M.F., tome 128, number 1 (2000), 103–149.
- [32] C. Taubes, *Casson’s invariant and gauge theory*. J. Differential Geom. 31 (1990), no. 2, 547–599.
- [33] C. Taubes and Y.J. Lee, *Periodic Floer homology and Seiberg-Witten-Floer cohomology*, J. Symplectic Geom. (2012), no. 1. 81–164.
- [34] K. Wehrheim, *Lagrangian boundary conditions for anti-self-dual instantons and the Atiyah-Floer conjecture*, J. Symplectic Geom. 3 (2005), no. 4, 703–747.
- [35] K. Wehrheim and C. Woodward, *Functoriality for Lagrangian correspondences in Floer theory*. Quantum Topol. 1 (2010), no. 2, 129–170.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824
E-mail address: `mhedden@math.msu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEVADA, RENO, RENO, NV 89557
E-mail address: `herald@unr.edu`

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405
E-mail address: `pkirk@indiana.edu`